

A Unifying Framework for Convergence Analysis of Approximate Newton Methods

Haishan Ye, Luo Luo

yhs12354123@gmail.com; rickyluoluo@gmail.com
Department of Computer Science and Engineering
Shanghai Jiao Tong University
800 Dong Chuan Road, Shanghai, China 200240

Zhihua Zhang

zhzhang@math.pku.edu.cn
School of Mathematical Sciences
Peking University
Beijing, China 100871

February 28, 2017

Abstract

Many machine learning models are reformulated as optimization problems. Thus, it is important to solve a large-scale optimization problem in big data applications. Recently, subsampled Newton methods have emerged to attract much attention for optimization due to their efficiency at each iteration, rectified a weakness in the ordinary Newton method of suffering a high cost in each iteration while commanding a high convergence rate. Other efficient stochastic second order methods are also proposed. However, the convergence properties of these methods are still not well understood. There are also several important gaps between the current convergence theory and the performance in real applications. In this paper, we aim to fill these gaps. We propose a unifying framework to analyze local convergence properties of second order methods. Based on this framework, our theoretical analysis matches the performance in real applications.

1. Introduction

Mathematical optimization is an importance pillar of machine learning. We consider the following optimization problem

$$\min_{x \in \mathbb{R}^d} F(x) = \frac{1}{n} \sum_{i=1}^n f_i(x), \quad (1)$$

where the $f_i(x)$ are smooth functions. Many machine learning models can be expressed as (1) where each f_i is the loss with respect to (w.r.t.) the i -th training sample. There are many examples such as logistic regressions, smoothed support vector machines, neural networks, and graphical models.

Many optimization algorithms to solve the problem in (1) are based on the following iteration:

$$x^{(t+1)} = x^{(t)} - \eta_t Q_t g(x^{(t)}), \quad t = 0, 1, 2, \dots,$$

where $\eta_t > 0$ is the step length. If Q_t is the identity matrix and $g(x^{(t)}) = \nabla F(x^{(t)})$, the resulting procedure is called *Gradient Descent* (GD) which achieves sublinear convergence for a general smooth convex objective function and linear convergence for a smooth-strongly convex objective function. When n is large, the full gradient method is inefficient due to its iteration cost scaling linearly in n . Consequently, stochastic gradient descent (SGD) has been a typical alternative [18, 12, 5]. In order to achieve cheaper cost in each iteration, such a method constructs an approximate gradient on a small mini-batch of data. However, the convergence rate can be significantly slower than that of the full gradient methods [15]. Thus, a great deal of efforts have been made to devise modification to achieve the convergence rate of the full gradient while keeping low iteration cost [10, 20, 21, 25].

If Q_t is a $d \times d$ positive definite matrix of containing the curvature information, this formulation leads us to *second-order* methods. It is well known that second order methods enjoy superior convergence rate in both theory and practice in contrast to *first-order* methods which only make use of the gradient information. The standard Newton method, where $Q_t = [\nabla^2 F(x^{(t)})]^{-1}$, $g(x^{(t)}) = \nabla F(x^{(t)})$ and $\eta_t = 1$, achieves a quadratic convergence rate for smooth-strongly convex objective functions. However, the *Newton method* takes $\mathcal{O}(nd^2 + d^3)$ cost per iteration, so it becomes extremely expensive when n or d is very large. As a result, one tries to construct an approximation of the Hessian in which way the update is computationally feasible, and while keeping sufficient second order information. One class of such methods are quasi-Newton methods, which are generalizations of the secant methods to find the root of the first derivative for multidimensional problems. The celebrated Broyden-Fletcher-Goldfarb-Shanno (BFGS) and its limited memory version (L-BFGS) are the most popular and widely used [16]. They take $\mathcal{O}(nd + d^2)$ cost per iteration.

Recently, when $n \gg d$, a class of called *subsampled Newton* methods have been proposed, which defines an approximate Hessian matrix with a small subset of samples. The most naive approach is to sample a subset of functions f_i randomly [19, 3, 24] to construct a subsampled Hessian. Erdogdu and Montanari [8] proposed a regularized subsampled Newton method called NewSamp. When the Hessian can be written as $\nabla^2 F(x) = [B(x)]^T B(x)$ where $B(x)$ is an available $n \times d$ matrix, Pilanci and Wainwright [17] used sketching techniques to approximate the Hessian and proposed *sketch Newton* method. Similarly, Xu et al. [24] proposed to sample rows of $B(x)$ with non-uniform probability distribution. Agarwal et al. [1] brought up an algorithm called LiSSA to approximate the inversion of Hessian directly.

Although the convergence performance of stochastic second order methods has been analyzed, the convergence properties are still not well understood. There are several important gaps lying between convergence theory and real application.

The first gap is the necessity of Lipschitz continuity of Hessian. In previous work, to achieve a linear-quadratic convergence rate, stochastic second order methods all assume that $\nabla^2 F(x)$ is Lipschitz continuous. However, in real application without this assumption, they might also converge to optimal point. For example, Erdogdu and Montanari [8] used NewSamp to successfully train smoothed-SVM in which case the Hessian is not Lipschitz continuous.

The second gap is about the sketched size of sketch Newton methods. To obtain a linear convergence, the sketched size is $\mathcal{O}(d\kappa^2)$ in [17] and then be improved to $\mathcal{O}(d\kappa)$ in [24] using Gaussian sketching matrices, where κ is the condition number of the Hessian matrix in question. However, the sketch Newton empirically performs well even when the Hessian matrix is ill-conditioned. Sketched size being several tens of times, or even several times of d can achieve a linear convergence rate in unconstrained optimization. But the theoretical result of Pilanci and Wainwright [17], Xu et al. [24] implies that sketched size may be beyond n in ill-condition cases.

The third gap is about the sample size in regularized subsampled Newton methods. In both [8] and [19], their theoretical analysis shows that the sample size of regularized subsampled Newton methods should be set as the same as the conventional subsampled Newton method. In practice, however, adding a large regularizer can obviously reduce the sample size while keeping convergence. Thus, this contradicts the extant theoretical analysis [8, 19].

In this paper, we aim to fill these gaps between the current theory and empirical performance. More specifically, we first cast these second order methods into an algorithmic framework that we call approximate Newton. Then we propose a general result for analysis of local convergence properties of second order methods. Based on this framework, we then give detailed theoretical analysis which matches the empirical performance. We summarize our contribution as follows:

- We propose a unifying framework (Theorem 3) to analyze local convergence properties of second order methods including stochastic and deterministic versions. The convergence performance of second order methods can be analyzed easily and systematically in this framework.
- We prove that the Lipschitz continuity condition of Hessian is not necessary for achieving linear and superlinear convergence in variants of subsampled Newton. But it is needed to obtain quadratic convergence. This explains the phenomenon that NewSamp [8] can be used to train smoothed SVM in which the Lipschitz continuity condition of Hessian is not satisfied. It also reveals the reason why previous stochastic second order methods, such as subsampled Newton, sketch Newton, LiSSA, etc., all achieve a linear-quadratic convergence rate.
- We prove that the sketched size is *independent* of the condition number of Hessian matrix which explains that sketched Newton performs well even when Hessian matrix is ill-conditioned.
- We provide a theoretical guarantee that adding a regularizer is an effective way to reduce sample size in subsampled Newton methods while keeping converging. Our theoretical analysis also shows that adding a regularizer will lead to poor convergence behavior because the sample size decreases.

1.1 Organization

The remainder of the paper is organized as follows. In Section 2 we present notation and preliminaries. In Section 3 we present a unifying framework for local convergence analysis

of second order methods. In Section 4 we analyze the local convergence properties of sketch Newton methods and prove that sketched size is independent of condition number of Hessian matrix. In section 5 we give the local convergence behaviors of several variants of subsampled Newton method. Especially, we reveal the relationship among sample size, regularizer and convergence rate. In Section 6, we derive the local convergence properties of inexact Newton method from our framework. In Section 7, we invalidate our theoretical results experimentally. Finally, we conclude our work in Section 8.

2. Notation and Preliminaries

Section 2.1 defines the notation used in this paper. Section 2.2 introduces matrices sketching techniques and their properties. Section 2.3 describes some important assumptions about object function.

2.1 Notation

Given a matrix $A = [a_{ij}] \in \mathbb{R}^{m \times n}$ of rank ℓ and a positive integer $k \leq \ell$, its condensed SVD is given as $A = U\Sigma V^T = U_k \Sigma_k V_k^T + U_{\setminus k} \Sigma_{\setminus k} V_{\setminus k}^T$, where U_k and $U_{\setminus k}$ contain the left singular vectors of A , V_k and $V_{\setminus k}$ contain the right singular vectors of A , and $\Sigma = \text{diag}(\sigma_1, \dots, \sigma_\ell)$ with $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_\ell > 0$ are the nonzero singular values of A . We will use $\sigma_{\max}(A)$ to denote the largest singular value and $\sigma_{\min}(A)$ to denote the smallest non-zero singular value. Thus, the condition number of A is defined by $\kappa(A) \triangleq \frac{\sigma_{\max}(A)}{\sigma_{\min}(A)}$. If A is positive semidefinite, then $U = V$ and the square root of A can be defined as $A^{1/2} = U\Sigma^{1/2}U^T$. It also holds that $\lambda_i(A) = \sigma_i(A)$, where $\lambda_i(A)$ is the i -th largest eigenvalue of A , $\lambda_{\max}(A) = \sigma_{\max}(A)$, and $\lambda_{\min}(A) = \sigma_{\min}(A)$.

Additionally, $\|A\|_F \triangleq (\sum_{i,j} a_{ij}^2)^{1/2} = (\sum_i \sigma_i^2)^{1/2}$ is the Frobenius norm of A and $\|A\| \triangleq \sigma_1$ is the spectral norm. Given a positive definite matrix M , $\|x\|_M \triangleq \|M^{1/2}x\|$ is called the M -norm of x . Give square matrices A and B with the same size, we denote $A \preceq B$ if $B - A$ is positive semidefinite.

2.2 Randomized sketching matrices

We first give an ϵ -subspace embedding property which will be used to sketch Hessian matrices. Then we list some useful types of randomized sketching matrices including Gaussian projection [9, 11], leverage score sampling [6], count sketch [4, 14, 13].

Definition 1 $S \in \mathbb{R}^{s \times m}$ is said to be an ϵ -subspace embedding matrix w.r.t. a fixed matrix $A \in \mathbb{R}^{m \times d}$ where $d < m$, if $\|SAx\|^2 = (1 \pm \epsilon)\|Ax\|^2$ (i.e., $(1 - \epsilon)\|Ax\|^2 \leq \|SAx\|^2 \leq (1 + \epsilon)\|Ax\|^2$) for all $x \in \mathbb{R}^d$.

From the definition of the ϵ -subspace embedding matrix, we can derive the following property directly.

Lemma 2 $S \in \mathbb{R}^{s \times m}$ is an ϵ -subspace embedding matrix w.r.t. the matrix $A \in \mathbb{R}^{m \times d}$ if and only if

$$(1 - \epsilon)A^T A \preceq A^T S^T S A \preceq (1 + \epsilon)A^T A.$$

Gaussian sketching matrix. The most classical sketching matrix is the Gaussian sketching matrix $S \in \mathbb{R}^{s \times m}$, whose entries are i.i.d. from the normal of mean 0 and variance $1/s$. Owing to the well-known concentration properties [23], Gaussian random matrices are very attractive. Besides, $s = \mathcal{O}(d/\epsilon^2)$ is enough to guarantee the ϵ -subspace embedding property for any fixed matrix $A \in \mathbb{R}^{m \times d}$. Moreover, $s = \mathcal{O}(d/\epsilon^2)$ is the tightest bound among known types of sketching matrices. However, the Gaussian random matrix is usually dense, so it is costly to compute SA .

Leverage score sketching matrix. A leverage score sketching matrix $S = D\Omega \in \mathbb{R}^{s \times m}$ w.r.t. $A \in \mathbb{R}^{m \times d}$ is defined by sampling probabilities p_i , a sampling matrix $\Omega \in \mathbb{R}^{m \times s}$ and a diagonal rescaling matrix $D \in \mathbb{R}^{s \times s}$. Specifically, we construct S as follows. For every $j = 1, \dots, s$, independently and with replacement, pick an index i from the set $\{1, 2, \dots, m\}$ with probability p_i , and set $\Omega_{ji} = 1$ and $\Omega_{jk} = 0$ for $k \neq i$ as well as $D_{jj} = 1/\sqrt{p_i s}$. The sampling probabilities p_i are the leverage scores of A defined as follows. Let $V \in \mathbb{R}^{m \times d}$ be the column orthonormal basis of A , and let $v_{i,*}$ denote the i -th row of V . Then $\ell_i \triangleq \|v_{i,*}\|^2/d$ for $i = 1, \dots, m$ are the leverage scores of A . To achieve an ϵ -subspace embedding property w.r.t. A , $s = \mathcal{O}(d \log d/\epsilon^2)$ is sufficient.

Sparse embedding matrix. A sparse embedding matrix $S \in \mathbb{R}^{s \times m}$ is such a matrix in each column of which there is only one nonzero entry uniformly sampled from $\{1, -1\}$ [4]. Hence, it is very efficient to compute SA , especially when A is sparse. To achieve an ϵ -subspace embedding property w.r.t. $A \in \mathbb{R}^{m \times d}$, $s = \mathcal{O}(d^2/\epsilon^2)$ is sufficient [13, 23].

Other sketching matrices such as Subsampled Randomized Hadamard Transformation [7, 9] as well as their properties can be found in the survey [23].

2.3 Assumptions and Notions

In this paper, we focus on the problem described in Eqn. (1). Moreover, we will make the following two assumptions.

Assumption 1 The objective function F is μ -strongly convex, that is,

$$F(y) \geq F(x) + [\nabla F(x)]^T(y - x) + \frac{\mu}{2}\|y - x\|^2, \text{ for } \mu > 0.$$

Assumption 2 $\nabla F(x)$ is L -Lipschitz continuous, that is,

$$\|\nabla F(x) - \nabla F(y)\| \leq L\|y - x\|, \text{ for } L > 0.$$

Assumptions 1 and 2 imply that for any $x \in \mathbb{R}^d$, we have

$$\mu I \preceq \nabla^2 F(x) \preceq LI,$$

where I is the identity matrix of appropriate size. With a little confusion, we define

$$\kappa \triangleq \frac{L}{\mu}.$$

In fact, κ is an upper bound of condition number of the Hessian matrix $\nabla^2 F(x)$ for any x .

Besides, if $\nabla^2 F(x)$ is Lipschitz continuous, then we have

$$\|\nabla^2 F(x) - \nabla^2 F(y)\| \leq \hat{L}\|x - y\|,$$

where $\hat{L} > 0$ is the Lipschitz constant of $\nabla^2 F(x)$.

Throughout this paper, we use notions of linear convergence rate, superlinear convergence rate and quadratic convergence rate. In our paper, the convergence rates we will use are defined w.r.t. $\|\cdot\|_{M^*}$, where $M^* = [\nabla^2 F(x^*)]^{-1}$ and x^* is the optimal solution to Problem (1). A sequence of vectors $\{x^{(t)}\}$ is said to converge linearly to a limit point x^* , if for some $0 < \rho < 1$,

$$\limsup_{t \rightarrow \infty} \frac{\|\nabla F(x^{(t+1)})\|_{M^*}}{\|\nabla F(x^{(t)})\|_{M^*}} = \rho.$$

Similarly, superlinear convergence and quadratic convergence are respectively defined as

$$\begin{aligned} \limsup_{t \rightarrow \infty} \frac{\|\nabla F(x^{(t+1)})\|_{M^*}}{\|\nabla F(x^{(t)})\|_{M^*}} &= 0, \\ \limsup_{t \rightarrow \infty} \frac{\|\nabla F(x^{(t+1)})\|_{M^*}}{\|\nabla F(x^{(t)})\|_{M^*}^2} &= \rho. \end{aligned}$$

We call it the linear-quadratic convergence rate if the following condition holds:

$$\|\nabla F(x^{(t+1)})\|_{M^*} \leq \rho_1 \|\nabla F(x^{(t)})\|_{M^*} + \rho_2 \|\nabla F(x^{(t)})\|_{M^*}^2,$$

where $0 < \rho_1 < 1$.

A small $\|\nabla F(x^{(t)})\|_{M^*}$ directly implies a small difference $\|x^{(t)} - x^*\|$ between $x^{(t)}$ and x^* . If $\|\nabla F(x^{(t)})\|_{M^*} \leq \epsilon$, then we have $\|\nabla F(x^{(t)})\| \leq \epsilon / \sqrt{\lambda_{\min}(M^*)} \leq \epsilon \sqrt{L}$ because $\|\nabla F(x^{(t)})\|_{M^*} = \|[M^*]^{1/2} \nabla F(x^{(t)})\| \geq \lambda_{\min}([M^*]^{1/2}) \|\nabla F(x^{(t)})\|$ and $\lambda_{\min}(M^*) \geq 1/L$. We also have $\|\nabla F(x^{(t)})\| = \|\nabla F(x^{(t)}) - \nabla F(x^*)\| \geq \mu \|x^{(t)} - x^*\|$ by the μ -strong convexity. Hence, we have

$$\|x^{(t)} - x^*\| \leq \frac{\sqrt{L}}{\mu} \epsilon$$

whenever $\|\nabla F(x^{(t)})\|_{M^*} \leq \epsilon$.

3. Approximate Newton Methods and Local Convergence Analysis

The existing variants of stochastic second order methods share some important attributes. First, these methods such as NewSamp [8], LiSSA [1], subsampled Newton with conjugate gradient [3], and subsampled Newton with non-uniformly sampling [24], all have the same convergence properties; that is, they have a linear-quadratic convergence rate.

Second, they also enjoy the same algorithm procedure summarized as follows. In each iteration, they first construct an approximate Hessian matrix $H^{(t)}$ such that

$$(1 - \epsilon_0)H^{(t)} \preceq \nabla^2 F(x^{(t)}) \preceq (1 + \epsilon_0)H^{(t)}, \quad (2)$$

where $0 \leq \epsilon_0 < 1$. Then they solve the following optimization problem

$$\min_p \frac{1}{2} p^T H^{(t)} p - p^T \nabla F(x^{(t)}) \quad (3)$$

approximately or exactly to obtain the direction vector $p^{(t)}$. Finally, their update equation is given as $x^{(t+1)} = x^{(t)} - p^{(t)}$. With this procedure, we regard these stochastic second order methods as *approximate Newton* methods.

In the following theorem, we propose a unifying framework which describes the convergence properties of the second order optimization procedure depicted above.

Theorem 3 *Let Assumption 1 and 2 hold. Suppose that $\nabla^2 F(x)$ exists and is continuous in a neighborhood of a minimizer x^* . $H^{(t)}$ is a positive definite matrix that satisfies Eqn. (2) with $0 \leq \epsilon_0 < 1$. Let $p^{(t)}$ be an approximate solution of Problem (3) such that*

$$\|\nabla F(x^{(t)}) - H^{(t)}p^{(t)}\| \leq \frac{\epsilon_1}{\kappa} \|\nabla F(x^{(t)})\|, \quad (4)$$

where $0 < \epsilon_1 < 1$. Consider the iteration $x^{(t+1)} = x^{(t)} - p^{(t)}$.

(a) *There exists a sufficient small value γ , $0 < \nu(t) < 1$, and $0 < \eta(t) < 1$ such that when $\|x^{(t)} - x^*\| \leq \gamma$, we have that*

$$\|\nabla F(x^{(t+1)})\|_{M^*} \leq \left(\epsilon_0 + \frac{\epsilon_1}{1 - \epsilon_0} + \frac{2\eta(t)}{1 - \epsilon_0} \right) \frac{1 + \nu(t)}{1 - \nu(t)} \|\nabla F(x^{(t)})\|_{M^*}. \quad (5)$$

Besides, $\nu(t)$ and $\eta(t)$ will go to 0 as $x^{(t)}$ goes to x^* .

(b) *Furthermore, if $\nabla^2 F(x)$ is Lipschitz continuous with parameter \hat{L} , and $x^{(t)}$ satisfies*

$$\|x^{(t)} - x^*\| \leq \frac{\mu}{\hat{L}\kappa} \nu(t), \quad (6)$$

where $0 < \nu(t) < 1$, then it holds that

$$\|\nabla F(x^{(t+1)})\|_{M^*} \leq \left(\epsilon_0 + \frac{\epsilon_1}{1 - \epsilon_0} \right) \frac{1 + \nu(t)}{1 - \nu(t)} \|\nabla F(x^{(t)})\|_{M^*} + \frac{2}{(1 - \epsilon_0)^2} \frac{\hat{L}\kappa}{\mu\sqrt{\mu}} \frac{(1 + \nu(t))^2}{1 - \nu(t)} \|\nabla F(x^{(t)})\|_{M^*}^2. \quad (7)$$

From Theorem 3, we can find some important insights. First, Theorem 3 provides sufficient conditions to get different convergence rates including super-linear and quadratic convergence rates. If $\left(\epsilon_0 + \frac{\epsilon_1}{1 - \epsilon_0} \right)$ is a constant, then sequence $\{x^{(t)}\}$ converges linearly because $\nu(t)$ and $\eta(t)$ will go to 0 as t goes to infinity. If we set $\epsilon_0 = \epsilon_0(t)$ and $\epsilon_1 = \epsilon_1(t)$ such that $\epsilon_0(t)$ and $\epsilon_1(t)$ decrease to 0 as t increases, then sequence $\{x^{(t)}\}$ will converge super-linearly. Similarly, if $\epsilon_0(t) = \mathcal{O}(\|\nabla F(x^{(t)})\|_{M^*})$, $\epsilon_1(t) = \mathcal{O}(\|\nabla F(x^{(t)})\|_{M^*})$, and $\nabla^2 F(x)$ is Lipschitz continuous, then sequence $\{x^{(t)}\}$ will converge quadratically.

Second, Theorem 3 makes it clear that the Lipschitz continuity of $\nabla^2 F(x)$ is *not necessary* for linear convergence and superlinear convergence of stochastic second order methods including Subsampled Newton method, Sketch Newton, NewSamp, etc. This reveals the reason why NewSamp can be used to train the smoothed SVM where the Lipschitz continuity of the Hessian matrix is not satisfied. The Lipschitz continuity condition is only needed to get a quadratic convergence or linear-quadratic convergence. This explains the phenomena that LiSSA[1], NewSamp [8], subsampled Newton with non-uniformly sampling [24], Sketched Newton [17] have linear-quadratic convergence rate because they all assume that the Hessian is Lipschitz continuous. In fact, it is well known that the Lipschitz continuity

Algorithm 1 Sketch Newton.

- 1: **Input:** $x^{(0)}$, $0 < \delta < 1$, $0 < \epsilon_0 < 1$;
 - 2: **for** $t = 0, 1, \dots$ until termination **do**
 - 3: Construct an ϵ_0 -subspace embedding matrix S for $B(x^{(t)})$ and where $\nabla^2 F(x)$ is of the form $\nabla^2 F(x) = (B(x^{(t)}))^T B(x^{(t)})$, and calculate $H^{(t)} = [B(x^{(t)})]^T S^T S B(x^{(t)})$;
 - 4: Calculate $p^{(t)} \approx \operatorname{argmin}_p \frac{1}{2} p^T H^{(t)} p - p^T \nabla F(x^{(t)})$;
 - 5: Update $x^{(t+1)} = x^{(t)} - p^{(t)}$;
 - 6: **end for**
-

condition of $\nabla^2 F(x)$ is not necessary to achieve a linear or superlinear convergence rate for inexact Newton methods.

Third, the unifying framework of Theorem 3 contains not only stochastic second order methods, but also the deterministic versions. For example, letting $H^{(t)} = \nabla^2 F(x^{(t)})$ and using conjugate gradient to get $p^{(t)}$, we obtain the famous “Newton-CG” method. In fact, different choice of $H^{(t)}$ and different way to calculate $p^{(t)}$ lead us to different second order methods. In the following sections, we will use this framework to analyze the local convergence performance of these second order methods in detail.

4. Sketch Newton Method

In this section, we use Theorem 3 to analyze the local convergence properties of Sketch Newton (Algorithm 1). We mainly focus on the case that the Hessian matrix is of the form

$$\nabla^2 F(x) = B(x)^T B(x) \quad (8)$$

where $B(x)$ is an explicitly available $n \times d$ matrix. Our result can be easily extended to the case that

$$\nabla^2 F(x) = B(x)^T B(x) + Q(x),$$

where $Q(x)$ is a positive semi-definite matrix related to the Hessian of regularizer.

Theorem 4 *Let $F(x)$ satisfy the conditions described in Theorem 3. Assume the Hessian matrix is given as Eqn. (8). Let $0 < \delta < 1$, $0 < \epsilon_0 < 1/2$ and $0 \leq \epsilon_1 < 1$ be given. $S \in \mathbb{R}^{\ell \times n}$ is an ϵ_0 -subspace embedding matrix w.r.t. $B(x)$ with probability at least $1 - \delta$, and direction vector $p^{(t)}$ satisfies Eqn. (4). Then Algorithm 1 has the following convergence properties:*

- (a) *There exists a sufficient small value γ , $0 < \nu(t) < 1$, and $0 < \eta(t) < 1$ such that when $\|x^{(t)} - x^*\| \leq \gamma$, then each iteration satisfies Eqn. (5) with probability at least $1 - \delta$.*
- (b) *If $\nabla^2 F(x^{(t)})$ is also Lipschitz continuous and $\{x^{(t)}\}$ satisfies Eqn. (6), then each iteration satisfies Eqn. (7) with probability at least $1 - \delta$.*

Theorem 4 directly provides a bound of the sketched size. Using the leverage score sketching matrix as an example, the sketched size $\ell = \mathcal{O}(d \log d / \epsilon_0^2)$ is sufficient. We compare our theoretical bound of the sketched size with the ones of Pilanci and Wainwright [17] and Xu et al. [24] in Table 1. As we can see, our sketched size is much smaller than the other two, especially when the Hessian matrix is ill-conditioned.

Table 1: Comparison with previous work

Reference	Sketched Size	Condition number free?
Pilanci and Wainwright [17]	$\mathcal{O}\left(\frac{d\kappa^2 \log d}{\epsilon_0^2}\right)$	No
Xu et al. [24]	$\mathcal{O}\left(\frac{d\kappa \log d}{\epsilon_0^2}\right)$	No
Our result (Theorem 4)	$\mathcal{O}\left(\frac{d \log d}{\epsilon_0^2}\right)$	Yes

Theorem 4 shows that the sketched size ℓ is *independent* on the condition number of the Hessian matrix $\nabla^2 F(x)$ just as shown in Table 1. This explains the phenomena that when the Hessian matrix is ill-conditioned, Sketch Newton performs well even when the sketched size is only several times of d . For a large condition number, the theoretical bounds of both Xu et al. [24] and Pilanci and Wainwright [17] may be beyond the number of samples n . Note that the theoretical results of [24] and [17] still hold in the constrained optimization problem. However, our result proves the effectiveness of the sketch Newton method for the unconstrained optimization problem in the ill-conditioned case.

Theorem 4 also contains the possibility of achieving an asymptotically super-linear rate by using an iteration-dependent sketching accuracy $\epsilon_0 = \epsilon_0(t)$. In particular, we present the following corollary.

Corollary 5 *$F(x)$ satisfies the the properties described in Theorem 3. Consider Algorithm 1 in which the iteration-dependent sketching accuracy is given as $\epsilon_0(t) = \frac{1}{\log(1+t)}$ and $p^{(t)} = [H^{(t)}]^{-1} \nabla F(x)$. If the initial point $x^{(0)}$ is close enough to the optimal point x^* , then sequence $\{x^{(t)}\}$ converges superlinearly.*

5. The Subsampled Newton method and Variants

In this section, we apply Theorem 3 to analyze subsampled Newton methods. First, we make the assumption that each $f_i(x)$ and $F(x)$ have the following properties:

$$\max_{1 \leq i \leq n} \|\nabla^2 f_i(x)\| \leq K < \infty, \quad (9)$$

$$\lambda_{\min}(\nabla^2 F(x)) \geq \sigma > 0. \quad (10)$$

It immediately follows from $\max_{i \leq n} \|\nabla^2 f_i(x)\| \leq K < \infty$ that $\|\nabla^2 F(x)\| \leq K$. Accordingly, if $\nabla^2 F(x)$ is ill-conditioned, then the value $\frac{K}{\sigma}$ is large.

5.1 The Subsampled Newton method

The Subsampled Newton method is depicted in Algorithm 2, and we now give its local convergence properties in the following theorem.

Theorem 6 *Let $F(x)$ satisfy the properties described in Theorem 3. Assume Eqn. (9) and Eqn. (10) hold and let $0 < \delta < 1$, $0 < \epsilon_0 < 1/2$ and $0 \leq \epsilon_1 < 1$ be given. $|\mathcal{S}|$ and $H^{(t)}$ are*

Algorithm 2 Subsampled Newton.

- 1: **Input:** $x^{(0)}$, $0 < \delta < 1$, $0 < \epsilon_0 < 1$;
 - 2: Set the sample size $|\mathcal{S}| \geq \frac{16K^2 \log(2d/\delta)}{\sigma^2 \epsilon_0^2}$.
 - 3: **for** $t = 0, 1, \dots$ until termination **do**
 - 4: Select a sample set \mathcal{S} , of size $|\mathcal{S}|$ and $H^{(t)} = \frac{1}{|\mathcal{S}|} \sum_{j \in \mathcal{S}} \nabla^2 f_j(x^{(t)})$;
 - 5: Calculate $p^{(t)} \approx \operatorname{argmin}_p \frac{1}{2} p^T H^{(t)} p - p^T \nabla F(x^{(t)})$;
 - 6: Update $x^{(t+1)} = x^{(t)} - p^{(t)}$;
 - 7: **end for**
-

set as in Algorithm 2, and the direction vector $p^{(t)}$ satisfies Eqn. (4). Then for $t = 1, \dots, T$, Algorithm 2 has the following convergence properties:

- (a) There exists a sufficient small value γ , $0 < \nu(t) < 1$, and $0 < \eta(t) < 1$ such that when $\|x^{(t)} - x^*\| \leq \gamma$, then each iteration satisfies Eqn. (5) with probability at least $1 - \delta$.
- (b) If $\nabla^2 F(x^{(t)})$ is also Lipschitz continuous with parameter \hat{L} and $\{x^{(t)}\}$ satisfies Eqn. (6), then each iteration satisfies Eqn. (7) with probability at least $1 - \delta$.

As we can see, Algorithm 2 almost has the same convergence properties as Algorithm 1 except several minor differences. The main difference is the construction manner of $H^{(t)}$ which should satisfy Eqn. (2). Algorithm 2 relies on the assumption that each $\|\nabla^2 f_i(x)\|$ is upper bounded (i.e., Eqn. (9) holds), while Algorithm 1 is built on the setting of the Hessian matrix as in Eqn. (8).

5.2 Regularized Subsampled Newton

In ill-conditioned cases (i.e., $\frac{K}{\sigma}$ is large), the subsampled Newton method in Algorithm 2 should take a lot of samples because the sample size $|\mathcal{S}|$ depends on $\frac{K}{\sigma}$ quadratically. To overcome this problem, one resorts to a regularized subsampled Newton method. The key idea is to add αI to the original subsampled Hessian just as described in Algorithm 3. Erdogdu and Montanari [8] proposed NewSamp which is another regularized subsampled Newton method depicted in Algorithm 4. In the following analysis, we prove that adding a regularizer is an effective way to reduce the sample size while keeping converging in theory.

We first give the theoretical analysis of local convergence properties of Algorithm 3.

Theorem 7 Let $F(x)$ satisfy the properties described in Theorem 3. Assume Eqns. (9) and (10) hold, and let $0 < \delta < 1$, $0 \leq \epsilon_1 < 1$ and $0 < \alpha$ be given. Assume β is a constant such that $0 < \beta < \alpha + \frac{\sigma}{2}$, the subsampled size $|\mathcal{S}|$ satisfies $|\mathcal{S}| \geq \frac{16K^2 \log(2d/\delta)}{\beta^2}$, and $H^{(t)}$ is constructed as in Algorithm 3. Define

$$\epsilon_0 = \max \left(\frac{\beta - \alpha}{\sigma + \alpha - \beta}, \frac{\alpha + \beta}{\sigma + \alpha + \beta} \right), \quad (11)$$

which implies that $0 < \epsilon_0 < 1$. Besides, the direction vector $p^{(t)}$ satisfies Eqn. (4). Then Algorithm 3 has the following convergence properties:

- (a) There exists a sufficient small value γ , $0 < \nu(t) < 1$, and $0 < \eta(t) < 1$ such that when $\|x^{(t)} - x^*\| \leq \gamma$, each iteration satisfies Eqn. (5) with probability at least $1 - \delta$.

(b) If $\nabla^2 F(x^{(t)})$ is also Lipschitz continuous with parameter \hat{L} and $\{x^{(t)}\}$ satisfies Eqn. (6), then each iteration satisfies Eqn. (7) with probability at least $1 - \delta$.

In Theorem 7 the parameter ϵ_0 mainly decides convergence properties of Algorithm 3. It is determined by two terms just as shown in Eqn. (11). These two terms depict the relationship among the sample size, regularizer αI , and convergence rate.

The first term describes the relationship between the regularizer αI and sample size. Without loss of generality, we set $\beta = \alpha$ which satisfies $0 < \beta < \alpha + \sigma/2$. Then the sample size $|\mathcal{S}| = \frac{16K^2 \log(2d/\delta)}{\alpha^2}$ decreases as α increases. Hence Theorem 7 gives a theoretical guarantee that adding the regularizer αI is an effective approach for reducing the sample size when K/σ is large. Conversely, if we want to sample a small part of f_i 's, then we should choose a large α . Otherwise, β will go to $\alpha + \sigma/2$ which means $\epsilon_0 = 1$, i.e., the sequence $\{x^{(t)}\}$ does not converge.

Though a large α can reduce the sample size, it is at the expense of slower convergence rate just as the second term shows. As we can see, $\frac{\alpha+\beta}{\sigma+\alpha+\beta}$ goes to 1 as α increases. Besides, ϵ_1 also has to decrease. Otherwise, $\epsilon_0 + \frac{\epsilon_1}{1-\epsilon_0}$ may be beyond 1 which means that Algorithm 3 will not converge.

In fact, slower convergence rate via adding a regularizer is because the sample size becomes small, which implies less curvature information is obtained. However, a small sample size implies low computational cost in each iteration. Therefore, a proper regularizer which balances the cost of each iteration and convergence rate is the key in the regularized subsampled Newton algorithm.

Algorithm 3 Regularized Subsample Newton.

```

1: Input:  $x^{(0)}$ ,  $0 < \delta < 1$ , regularizer parameter  $\alpha$ , sample size  $|\mathcal{S}|$ ;
2: for  $t = 0, 1, \dots$  until termination do
3:   Select a sample set  $\mathcal{S}$ , of size  $|\mathcal{S}|$  and  $H^{(t)} = \frac{1}{|\mathcal{S}|} \sum_{j \in \mathcal{S}} \nabla^2 f_j(x^{(t)}) + \alpha I$ ;
4:   Calculate  $p^{(t)} \approx \operatorname{argmin}_p \frac{1}{2} p^T H^{(t)} p - p^T \nabla F(x^{(t)})$ 
5:   Update  $x^{(t+1)} = x^{(t)} - p^{(t)}$ ;
6: end for

```

Algorithm 4 NewSamp.

```

1: Input:  $x^{(0)}$ ,  $0 < \delta < 1$ ,  $r$ , sample size  $|\mathcal{S}|$ ;
2: for  $t = 0, 1, \dots$  until termination do
3:   Select a sample set  $\mathcal{S}$ , of size  $|\mathcal{S}|$  and get  $H_{|\mathcal{S}|}^{(t)} = \frac{1}{|\mathcal{S}|} \sum_{j \in \mathcal{S}} \nabla^2 f_j(x^{(t)})$ ;
4:   Compute rank  $r + 1$  truncated SVD deompostion of  $H_{|\mathcal{S}|}^{(t)}$  to get  $U_{r+1}$  and  $\hat{\Lambda}_{r+1}$ . Construct
       $H^{(t)} = H_{|\mathcal{S}|}^{(t)} + U_{\setminus r}(\hat{\Lambda}_{r+1}^{(t)} I - \hat{\Lambda}_{\setminus r})U_{\setminus r}^T$ 
5:   Calculate  $p^{(t)} \approx \operatorname{argmin}_p \frac{1}{2} p^T H^{(t)} p - p^T \nabla F(x^{(t)})$ 
6:   Update  $x^{(t+1)} = x^{(t)} - p^{(t)}$ ;
7: end for

```

Next, we give the theoretical analysis of local convergence properties of NewSamp (Algorithm 4).

Theorem 8 *Let $F(x)$ satisfy the properties described in Theorem 3. Assume Eqn. (9) and Eqn. (10) hold and let $0 < \delta < 1$ and target rank r be given. Let β be a constant such that $0 < \beta < \frac{\lambda_{r+1}^{(t)}}{2}$, where $\lambda_{r+1}^{(t)}$ is the $(r+1)$ -th eigenvalue of $\nabla^2 F(x^{(t)})$. Set the subsampled size $|\mathcal{S}|$ such that $|\mathcal{S}| \geq \frac{16K^2 \log(2d/\delta)}{\beta^2}$, and define*

$$\epsilon_0 = \max \left(\frac{\beta}{\lambda_{r+1}^{(t)} - \beta}, \frac{2\beta + \lambda_{r+1}^{(t)}}{\sigma + 2\beta + \lambda_{r+1}^{(t)}} \right), \quad (12)$$

which implies $0 < \epsilon_0 < 1$. Assume the direction vector $p^{(t)}$ satisfies Eqn. (4). Then for $t = 1, \dots, T$, Algorithm 4 has the following convergence properties:

- (a) *There exists a sufficient small value γ , $0 < \nu(t) < 1$, and $0 < \eta(t) < 1$ such that when $\|x^{(t)} - x^*\| \leq \gamma$, each iteration satisfies Eqn. (5) with probability at least $1 - \delta$.*
- (b) *If $\nabla^2 F(x^{(t)})$ is also Lipschitz continuous with parameter \hat{L} and $\{x^{(t)}\}$ satisfies Eqn. (6), then each iteration satisfies Eqn. (7) with probability at least $1 - \delta$.*

Similar to Theorem 7, parameter ϵ_0 in NewSamp is also determined by two terms. The first term reveals the relationship between the target rank r and sample size. Without loss of generality, we can set $\beta = \lambda_{r+1}^{(t)}/4$. Then the sample size is linear to $1/[\lambda_{r+1}^{(t)}]^2$. Hence, a small r means that a small sample size is sufficient. Conversely, if we want to sample a small portion of f_i 's, then we should choose a small r . Otherwise, β will go to $\lambda_{r+1}^{(t)}/2$ which means $\epsilon_0 = 1$, i.e., the sequence $\{x^{(t)}\}$ does not converge. The second term shows that a small sample size will lead to a poor convergence rate. If we set $r = 0$ and $\beta = \lambda_1/2$, then ϵ_0 will be $1 - \frac{1}{1+2\lambda_1/\sigma}$. Consequently, the convergence rate of NewSamp is almost the same as gradient descent. Similar to Algorithm 3, a small r means a precise solution to Problem (3) and the initial point $x^{(0)}$ being close to the optimal point x^* .

It is worth pointing out that Theorem 8 explains the empirical results that NewSamp is applicable in training SVM in which the Lipschitz continuity condition of $\nabla^2 F(x)$ is not satisfied [8].

We now conduct comparison between Theorem 7 and Theorem 8. We mainly focus on the parameter ϵ_0 in these two theorems which mainly determines convergence properties of Algorithm 3 and Algorithm 4. Specifically, if we set $\alpha = \beta + \lambda_{r+1}^{(t)}$ in Eqn. (11), then $\epsilon_0 = \frac{2\beta + \lambda_{r+1}^{(t)}}{\sigma + 2\beta + \lambda_{r+1}^{(t)}}$ which equals to the second term on the right-hand side in Eqn. (12). Hence, we can regard NewSamp as a special case of Algorithm 3. However, NewSamp provides an approach for automatical choice of α .

Recall that NewSamp includes another parameter: the target rank r . Thus, NewSamp and Algorithm 3 have the same number of free parameters. If r is not properly chosen, NewSamp will still have poor performance. Therefore, Algorithm 3 is theoretically preferred because NewSamp needs extra cost to perform SVDs.

5.3 Subsampled Hessian and Gradient

In fact, we can also subsample gradient to accelerate the subsampled Newton method. The detailed procedure is presented in Algorithm 5 [3, 19].

Algorithm 5 Subsampled Hessian and Subsampled Gradient.

```

1: Input:  $x^{(0)}$ ,  $0 < \delta < 1$ ,  $0 < \epsilon_0 < 1$ ;
2: Set the sample size  $|\mathcal{S}_H|$  and  $|\mathcal{S}_g|$ .
3: for  $t = 0, 1, \dots$  until termination do
4:   Select a sample set  $\mathcal{S}_H$ , of size  $|\mathcal{S}|$  and construct  $H^{(t)} = \frac{1}{|\mathcal{S}|} \sum_{j \in \mathcal{S}} \nabla^2 f_j(x^{(t)})$ ;
5:   Select a sample set  $\mathcal{S}_g$  of size  $|\mathcal{S}_g|$  and calculate  $g(x^{(t)}) = \frac{1}{|\mathcal{S}_g|} \sum_{i \in \mathcal{S}_g} \nabla f_i(x^{(t)})$ .
6:   Calculate  $p^{(t)} = [H^{(t)}]^{-1} g(x^{(t)})$ ;
7:   Update  $x^{(t+1)} = x^{(t)} - p^{(t)}$ ;
8: end for

```

Theorem 9 *Let $F(x)$ satisfy the properties described in Theorem 3. We also assume Eqn. (9) and Eqn. (10) hold and let $0 < \delta < 1$ and $0 < \epsilon_0 < 1/2$ be given. Let $|\mathcal{S}_H|$ and $|\mathcal{S}_g|$ be set such that Eqn. (2) holds and it holds that*

$$\|g(x^{(t)}) - \nabla F(x^{(t)})\| \leq \frac{\epsilon_2}{\kappa} \|\nabla F(x^{(t)})\|.$$

The direction vector $p^{(t)}$ is computed as in Algorithm 5. Then for $t = 1, \dots, T$, we have the following convergence properties:

- (a) *There exists a sufficient small value γ , $0 < \nu(t) < 1$, and $0 < \eta(t) < 1$ such that when $\|x^{(t)} - x^*\| \leq \gamma$, then for each iteration, it holds that*

$$\|\nabla F(x^{(t+1)})\|_{M^*} \leq (\epsilon_0 + 2\epsilon_2 + 4\eta(t)) \frac{1 + \nu(t)}{1 - \nu(t)} \|\nabla F(x^{(t)})\|_{M^*}$$

with probability at least $1 - \delta$.

- (b) *If $\nabla^2 F(x^{(t)})$ is also Lipschitz continuous with parameter \hat{L} and $\{x^{(t)}\}$ satisfies Eqn. (6), then for each iteration, it holds that*

$$\|\nabla F(x^{(t+1)})\|_{M^*} \leq (\epsilon_0 + 2\epsilon_2) \frac{1 + \nu(t)}{1 - \nu(t)} \|\nabla F(x^{(t)})\|_{M^*} + \frac{8\hat{L}\kappa}{\mu\sqrt{\mu}} \frac{(1 + \nu(t))^2}{1 - \nu(t)} \|\nabla F(x^{(t)})\|_{M^*}^2.$$

with probability at least $1 - \delta$.

In common cases, subsampled gradient $g(x^{(t)})$ needs to subsample over 80% of samples to guarantee convergence of the algorithm. Roosta-Khorasani and Mahoney [19] showed that it needs $|\mathcal{S}_g| \geq G(x^{(t)})^2 \kappa^2 / (\nu^2(t) \|\nabla F(x^{(t)})\|^2)$, where $G(x^{(t)}) = \max_i \|\nabla f_i(x^{(t)})\|$ for $i = 1, \dots, n$. When $x^{(t)}$ is close to x^* , $\|\nabla F(x^{(t)})\|$ is close to 0. Hence $|\mathcal{S}_g|$ will go to n as iteration goes. This is the reason why the Newton method and variants of the subsampled Newton method are very sensitive to the accuracy of subsampled gradient.

6. Inexact Newton Methods

Let $H^{(t)} = \nabla^2 F(x^{(t)})$, that is $\epsilon_0 = 0$. Then Theorem 3 depicts the convergence properties of inexact Newton methods.

Theorem 10 Let $F(x)$ satisfy the properties described in Theorem 3, and $p^{(t)}$ be a direction vector such that

$$\|\nabla F(x^{(t)}) - \nabla^2 F(x^{(t)})p^{(t)}\| \leq \frac{\epsilon_1}{\kappa} \|\nabla F(x^{(t)})\|,$$

where $0 < \epsilon_1 < 1$ is a constant. Consider the iteration $x^{(t+1)} = x^{(t)} - p^{(t)}$.

(a) There exists a sufficient small value γ , $0 < \nu(t) < 1$, and $0 < \eta(t) < 1$ such that when $\|x^{(t)} - x^*\| \leq \gamma$, then it holds that

$$\|\nabla F(x^{(t+1)})\|_{M^*} \leq \left(\epsilon_0 + \frac{\epsilon_2}{1 - \epsilon_0} + \frac{2\eta(t)}{1 - \epsilon_0} \right) \frac{1 + \nu(t)}{1 - \nu(t)} \|\nabla F(x^{(t)})\|_{M^*}.$$

(b) If $\nabla^2 F(x)$ is also Lipschitz continuous with parameter \hat{L} , and $\{x^{(t)}\}$ satisfies Eqn. (6), then it holds that

$$\|\nabla F(x^{(t+1)})\|_{M^*} \leq \left(\epsilon_0 + \frac{\epsilon_2}{1 - \epsilon_0} \right) \frac{1 + \nu(t)}{1 - \nu(t)} \|\nabla F(x^{(t)})\|_{M^*} + \frac{2\hat{L}\kappa}{(1 - \epsilon_0)^2 \mu \sqrt{\mu}} \frac{(1 + \nu(t))^2}{1 - \nu(t)} \|\nabla F(x^{(t)})\|_{M^*}^2.$$

6.1 LiSSA

LiSSA aims to approximate the inverse of Hessian matrix stochastically [1]. This is equivalent to solving the following linear equation

$$\nabla^2 F(x^{(t)})p = \nabla F(x^{(t)})$$

inexactly by a stochastic method. Thus, LiSSA can be regarded as a special case of inexact Newton method. Similarly, LiSSA-Sample also belongs to inexact Newton methods but with different way to solve the above linear equation. A detailed comparison between the subsampled Newton method and LiSSA can be found in [2].

7. Empirical Analysis

In this section, we validate our theoretical results about unnecessary of the Lipschitz continuity condition of $\nabla^2 F(x)$, sketched size of the sketch Newton, and sample size of regularized Newton, experimentally.

Table 2: Datasets Description

Dataset	n	d	source
mushrooms	8124	112	UCI
a9a	32561	123	UCI
Coverttype	581012	54	UCI

7.1 Unnecessity of Lipschitz continuity of Hessian

We conduct experiment on the primal problem for the linear SVM which can be written as

$$\min_x F(x) = \frac{1}{2} \|x\|^2 + \frac{C}{2n} \sum_{i=1}^n \ell(b_i, \langle x, a_i \rangle)$$

where (a_i, b_i) denotes the training data, x defines the separating hyperplane, $C > 0$, and $\ell(\cdot)$ is the loss function. In our experiment, we choose Hinge-2 loss as our loss function whose definition is

$$\ell(b, \langle x, a \rangle) = \max(0, 1 - b\langle x, a \rangle)^2.$$

Let $SV^{(t)}$ denote the set of indices of all the support vectors at iteration t , i.e.,

$$SV^{(t)} = \{i : b_i \langle x^{(t)}, a_i \rangle < 1\}.$$

Then the Hessian matrix of $F(x^{(t)})$ can be written as

$$\nabla^2 F(x^{(t)}) = I + \frac{1}{n} \sum_{i \in SV^{(t)}} a_i a_i^T.$$

From the above equation, we can see that $\nabla^2 F(x)$ is not Lipschitz continuous.

Without loss of generality, we use the Subsampled Newton method (Algorithm 2) in our experiment. We sample 5% support vectors in each iteration. Our experiments on three datasets whose detailed description is in Table 2 and report our results in Figure 1.

From Figure 1, we can see that Subsampled Newton converges linearly and the Newton method converges superlinearly. This matches our theory that the Lipschitz continuity of $\nabla^2 F(x)$ is not necessary to achieve a linear or superlinear convergence rate.




Figure 1: Convergence properties on different datasets.

7.2 Sketched Size of Sketch Newton

Now we invalidate that our theoretical result that sketched size is independent of the condition number of the Hessian matrix in Sketch Newton. To control the condition number of the Hessian conveniently, we conduct the experiment on least squares regression which is defined as

$$\min_x \frac{1}{2} \|Ax - b\|^2. \quad (13)$$

In each iteration, the Hessian matrix is $A^T A$. In our experiment, A is a 10000×54 matrix. And we set the singular values σ_i of A as:

$$\sigma_i = 1.2^{-i}.$$

Then the condition number of A is $\kappa(A) = 1.2^{54} = 1.8741 \times 10^4$. We use different sketch matrices in Sketch Newton (Algorithm 1) and set different values of the sketched size ℓ . We report our empirical results in Figure 2.

From Figure 2, we can see that Sketch Newton performs well when the sketch size ℓ is several times of d for all different sketching matrices. Moreover, the corresponding algorithms converge linearly. This matches our theory that sketched size is independent of the condition number of Hessian matrix to achieve a linear convergence rate. In contrast, the theoretical result of [24] shows that sketched size is $\ell = d * \kappa(A) = 1.02 \times 10^6$ bigger than $n = 10^4$.




Figure 2: Convergence properties of different sketched sizes

7.3 Sample Size of Regularized Subsampled Newton

We also choose least squares regression defined in Eqn. (13) in our experiment to validate the theory that adding a regularizer is an effective approach to reducing the sample size while keeping convergence in Subsampled Newton. Let $A \in \mathbb{R}^{n \times d}$ where $n = 8000$ and $d = 5000$. Hence Sketch Newton can not be used in this case because n and d are close to each other. In our experiment, we set different sample sizes $|\mathcal{S}|$. For each $|\mathcal{S}|$ we choose different regularizer terms α and different target ranks r . We report our results in Figures 3 and 4.

As we can see, if the sample size $|\mathcal{S}|$ is small, then we should choose a large α in Algorithm 3; otherwise, the algorithm will diverge. However, if the regularizer term α is too large, then the algorithm will converge slowly. Besides, increasing the sample size and choosing a proper regularizer will improve convergence properties obviously. When $|\mathcal{S}| = 600$, it only needs about 1200 iterations to obtain a precise solution while it needs about 8000 iterations when $|\mathcal{S}| = 100$. Similarly, if the sample size $|\mathcal{S}|$ is small, then we should choose a small target rank in NewSamp. Otherwise NewSamp may diverge. Also if the target rank is not chosen properly, then NewSamp will have poor convergence properties. Furthermore, comparing Figures 3 and 4, we can see that the two algorithms have similar

convergence properties. This invalidates the theoretical result that NewSamp provides a method to choose α automatically. Our empirical analysis matches the theoretical analysis in Subsection 5.2 very well.




Figure 3: Convergence properties of Regularized Subsampled Newton




Figure 4: Convergence properties of NewSamp

8. Conclusion

In this paper, we have proposed a framework to analyze the local convergence properties of second order methods including stochastic and deterministic versions. This framework reveals some important convergence properties of the subsampled Newton method and sketch Newton method, which are unknown before. The most important thing is our analysis lay the theoretical foundation of several important stochastic second order methods.

We believe that this framework might also provide some useful insights for developing new subsampled Newton-type algorithms. We would like to address this issue in future.

Appendix A. Some Important Lemmas

In this section, we give several important lemmas which will be used in the proof of the theorems of this paper.

Lemma 11 *If A and B are $d \times d$ symmetric positive matrices, and $(1-\epsilon_0)B \preceq A \preceq (1+\epsilon_0)B$ where $0 < \epsilon_0 < 1$, then we have*

$$\|A^{1/2}B^{-1}A^{1/2} - I\| \leq \epsilon_0,$$

where I is the identity matrix.

Proof Because $A \preceq (1+\epsilon_0)B$, we have $z^T[A - (1+\epsilon_0)B]z \leq 0$ for any nonzero $z \in \mathbb{R}^d$. This implies $\frac{z^T A z}{z^T B z} \leq 1 + \epsilon_0$ for any $z \neq 0$. Subsequently,

$$\begin{aligned} \lambda_{\max}(B^{-1}A) &= \lambda_{\max}(B^{-1/2}AB^{-1/2}) \\ &= \max_{u \neq 0} \frac{u^T B^{-1/2}AB^{-1/2}u}{u^T u} \\ &= \max_{z \neq 0} \frac{z^T A z}{z^T B z} \\ &\leq 1 + \epsilon_0, \end{aligned}$$

where the last equality is obtained by setting $z = B^{-1/2}u$. Similarly, we have $\lambda_{\min}(B^{-1}A) \geq 1 - \epsilon_0$. Since $B^{-1}A$ and $A^{1/2}B^{-1}A^{1/2}$ are similar, the eigenvalues of $A^{1/2}B^{-1}A^{1/2}$ are all between $1 - \epsilon_0$ and $1 + \epsilon_0$. Therefore, we have

$$\|A^{1/2}B^{-1}A^{1/2} - I\| \leq \epsilon_0.$$

■

Lemma 12 ([22]) *Let X_1, X_2, \dots, X_k be independent, random, symmetric, real matrices of size $d \times d$ with $0 \preceq X_i \preceq LI$, where I is the $d \times d$ identity matrix. Let $Y = \sum_{i=1}^k X_i$, $\mu_{\min} = \lambda_{\min}(\mathbb{E}[Y])$ and $\mu_{\max} = \lambda_{\max}(\mathbb{E}[Y])$. Then,*

$$\mathbb{P}(\lambda_{\min}(Y) \leq (1 - \epsilon)\mu_{\min}) \leq d \cdot e^{-\epsilon^2 \mu_{\min}/L}$$

Lemma 13 ([22]) *Given a matrix $A \in \mathbb{R}^{m \times n}$, construct an $m \times n$ random matrix R such that*

$$\mathbb{E}[R] = A \quad \text{and} \quad \|R\| \leq L.$$

Compute the per-sample second moment:

$$M = \max\{\|\mathbb{E}[RR^T]\|, \|\mathbb{E}[R^T R]\|\}.$$

Form the matrix sampling estimator

$$\bar{R} = \frac{1}{s} \sum_{i=1}^s R_i, \text{ where each } R_i \text{ is an independent copy of } R.$$

Then, for all $t \geq 0$

$$\mathbb{P} [\|\bar{R} - A\| \geq t] \leq (m + n) \exp \left(\frac{-st^2/2}{M + 2Lt/3} \right).$$

Lemma 14 Assume (9) and (10) hold. Let $0 < \delta < 1$, $0 < \epsilon < 1$ and $0 < c$ be given. If we sample f_i 's uniformly with the sample size $|\mathcal{S}|$ and construct $H^{(t)} = \frac{1}{|\mathcal{S}|} \sum_{j \in \mathcal{S}} \nabla^2 f_j(x^{(t)})$, then we have the following results:

(a) If $|\mathcal{S}| \geq \frac{16K^2 \log(2d/\delta)}{c^2 \epsilon^2}$, it holds that

$$\|H^{(t)} - \nabla^2 F(x^{(t)})\| \leq \epsilon c.$$

(b) If $|\mathcal{S}| \geq \frac{K \log(2d/\delta)}{\sigma \epsilon^2}$, it holds that

$$\lambda_{\min}(H^{(t)}) \geq (1 - \epsilon)\sigma.$$

Proof Consider $|\mathcal{S}|$ i.i.d random matrices $H_j^{(t)}, j = 1, \dots, |\mathcal{S}|$ such that $\mathbb{P}(H_j^{(t)} = \nabla^2 f_i(x^{(t)})) = 1/n$ for all $i = 1, \dots, n$. Then, we have $\mathbb{E}(H_j^{(t)}) = \nabla^2 F(x^{(t)})$ for all $j = 1, \dots, |\mathcal{S}|$. By (9) and the positive semi-definite property of $H_j^{(t)}$, we have $\lambda_{\max}(H_j^{(t)}) \leq K$ and $\lambda_{\min}(H_j^{(t)}) \geq 0$. By Lemma 12, we have that if $|\mathcal{S}| \geq \frac{K \log(d/\delta)}{\sigma \epsilon^2}$, $\lambda_{\min}(H^{(t)}) \geq (1 - \epsilon)\sigma$ holds with probability at least $1 - \delta$.

We define random matrices $X_j = H_j^{(t)} - \nabla^2 F(x^{(t)})$ for all $j = 1, \dots, |\mathcal{S}|$. We have $\mathbb{E}[X_j] = 0$, $\|X_j\| \leq 2K$ and $\|X_j\|^2 \leq 4K^2$. By Lemma 13, we have

$$\mathbb{P}(\|H^{(t)} - \nabla^2 F(x^{(t)})\| \geq \epsilon c) \leq 2d \exp^{-\frac{c^2 \epsilon^2 |\mathcal{S}|}{16K^2}}.$$

When $|\mathcal{S}| \geq \frac{16K^2 \log(2d/\delta)}{c^2 \epsilon^2}$, $\|H^{(t)} - \nabla^2 F(x^{(t)})\| \leq \epsilon c$ holds with probability at least $1 - \delta$. ■

Appendix B. Proofs of theorems of Section 3

Proof of Theorem 3 By Assumption 1 and 2, we have that $F(x)$ is μ -strongly convex and $\nabla F(x)$ is L -Lipschitz continuous. Hence, we have

$$\mu \leq \lambda_{\min}(\nabla^2 F(x)) \leq \lambda_{\max}(\nabla^2 F(x)) \leq L.$$

Hence, for any x in domain, it holds that

$$\kappa = \frac{L}{\mu} \geq \kappa(\nabla^2 F(x)).$$

By Taylor's theorem, we obtain

$$\begin{aligned}
& \nabla F(x^{(t+1)}) \\
&= \nabla F(x^{(t)}) + \nabla^2 F(x^{(t)})(-p^{(t)}) + \int_0^1 [\nabla^2 F(x^{(t)} + sp^{(t)}) - \nabla^2 F(x^{(t)})](-p^{(t)})ds \\
&= \nabla F(x^{(t)}) - \nabla^2 F(x^{(t)})[H^{(t)}]^{-1}\nabla F(x^{(t)}) + \nabla^2 F(x^{(t)})[H^{(t)}]^{-1}\nabla F(x^{(t)}) - \nabla^2 F(x^{(t)})p^{(t)} \\
&\quad + \int_0^1 [\nabla^2 F(x^{(t)} + sp^{(t)}) - \nabla^2 F(x^{(t)})](-p^{(t)})ds \\
&= [\nabla^2 F(x^{(t)})]^{\frac{1}{2}} \left(I - [\nabla^2 F(x^{(t)})]^{\frac{1}{2}}[H^{(t)}]^{-1}[\nabla^2 F(x^{(t)})]^{\frac{1}{2}} \right) [\nabla^2 F(x^{(t)})]^{-\frac{1}{2}} \nabla F(x^{(t)}) \\
&\quad + \nabla^2 F(x^{(t)})([H^{(t)}]^{-1}\nabla F(x^{(t)}) - p^{(t)}) + \int_0^1 [\nabla^2 F(x^{(t)} + sp^{(t)}) - \nabla^2 F(x^{(t)})](-p^{(t)})ds.
\end{aligned}$$

Hence, we have the following identity

$$\begin{aligned}
[\nabla^2 F(x^{(t)})]^{-\frac{1}{2}} \nabla F(x^{(t+1)}) &= \left(I - [\nabla^2 F(x^{(t)})]^{\frac{1}{2}}[H^{(t)}]^{-1}[\nabla^2 F(x^{(t)})]^{\frac{1}{2}} \right) [\nabla^2 F(x^{(t)})]^{-\frac{1}{2}} \nabla F(x^{(t)}) \\
&\quad + [\nabla^2 F(x^{(t)})]^{\frac{1}{2}}([H^{(t)}]^{-1}\nabla F(x^{(t)}) - p^{(t)}) \\
&\quad + [\nabla^2 F(x^{(t)})]^{-\frac{1}{2}} \int_0^1 [\nabla^2 F(x^{(t)} + sp^{(t)}) - \nabla^2 F(x^{(t)})](-p^{(t)})ds.
\end{aligned}$$

For notational simplicity, we denote $M = [\nabla^2 F(x^{(t)})]^{-1}$ and $M^* = [\nabla^2 F(x^*)]^{-1}$. Then we can obtain

$$\begin{aligned}
\|\nabla F(x^{(t+1)})\|_M &\leq \left\| I - [\nabla^2 F(x^{(t)})]^{\frac{1}{2}}[H^{(t)}]^{-1}[\nabla^2 F(x^{(t)})]^{\frac{1}{2}} \right\| \|\nabla F(x^{(t)})\|_M \\
&\quad + \|[\nabla^2 F(x^{(t)})]^{\frac{1}{2}}\| \| [H^{(t)}]^{-1}\nabla F(x^{(t)}) - p^{(t)} \| \\
&\quad + \|[\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}\| \int_0^1 \|\nabla^2 F(x^{(t)} + sp^{(t)}) - \nabla^2 F(x^{(t)})\| \|p^{(t)}\| ds.
\end{aligned}$$

We bound the three terms on the right-hand side of the above equation respectively.

For the first term, using Lemma 11, we have

$$\left\| I - [\nabla^2 F(x^{(t)})]^{\frac{1}{2}}[H^{(t)}]^{-1}[\nabla^2 F(x^{(t)})]^{\frac{1}{2}} \right\| \|\nabla F(x^{(t)})\|_M \leq \epsilon_0 \|\nabla F(x^{(t)})\|_M.$$

For the second term, by the fact that $\|AB\| \geq \|A\|\sigma_{\min}(B)$ and condition

$$\|\nabla F(x^{(t)}) - H^{(t)}p^{(t)}\| \leq \frac{\epsilon_1}{\kappa} \|\nabla F(x^{(t)})\| \leq \frac{\epsilon_1}{\kappa(\nabla^2 F(x^{(t)}))} \|\nabla F(x^{(t)})\|,$$

we obtain

$$\begin{aligned}
& \|[\nabla^2 F(x^{(t)})]^{\frac{1}{2}}\| \| [H^{(t)}]^{-1} \nabla F(x^{(t)}) - p^{(t)} \| \\
&= \frac{\|[\nabla^2 F(x^{(t)})]^{\frac{1}{2}}\|}{\lambda_{\min}([\nabla^2 F(x^{(t)})]^{-\frac{1}{2}})} \lambda_{\min}([\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}) \| [H^{(t)}]^{-1} \| \|\nabla F(x^{(t)}) - H^{(t)} p^{(t)}\| \\
&\leq \frac{\epsilon_1}{\kappa(\nabla^2 F(x^{(t)}))} \frac{\|[\nabla^2 F(x^{(t)})]^{\frac{1}{2}}\|}{\lambda_{\min}([\nabla^2 F(x^{(t)})]^{-\frac{1}{2}})} \| [H^{(t)}]^{-1} \| (\lambda_{\min}([\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}) \|\nabla F(x^{(t)})\|) \\
&\leq \frac{\epsilon_1}{\kappa(\nabla^2 F(x^{(t)}))} \|\nabla^2 F(x^{(t)})\| \| [H^{(t)}]^{-1} \| \|\nabla F(x^{(t)})\|_M \\
&\leq \frac{\epsilon_1}{1 - \epsilon_0} \|\nabla F(x^{(t)})\|_M.
\end{aligned}$$

For the third term, we bound it for the case that $\nabla^2 F(x)$ is not Lipschitz continuous and the case $\nabla^2 F(x)$ is Lipschitz continuous respectively.

First, we consider the case that $\nabla^2 F(x)$ is not Lipschitz continuous but is continuous close to the optimal point x^* . Because $\nabla^2 F(x)$ is continuous near x^* , there exists a sufficient small value γ such that it holds that

$$\|[\nabla^2 F(x^*)]^{-1} - [\nabla^2 F(x^{(t)})]^{-1}\| < \frac{\nu(t)}{L}, \quad (14)$$

and

$$\|\nabla^2 F(x^*) - \nabla^2 F(x^{(t)})\| < \frac{\eta(t)\mu}{\sqrt{\kappa}}, \quad (15)$$

when $\|x^{(t)} - x^*\| \leq \gamma$. Therefore, $\nu(t)$ and $\eta(t)$ will go to 0 as $x^{(t)}$ goes to x^* .

By μ -strong convexity, we have $\|[\nabla^2 F(x^{(t)})]^{-1}\| \leq \frac{1}{\mu}$ for all $x^{(t)}$ sufficiently close to x^* . Because of Eqn. (2), we have

$$\|[H^{(t)}]^{-1}\| \leq (1 + \epsilon_0) \|[\nabla^2 F(x^{(t)})]^{-1}\| \leq \frac{1}{(1 - \epsilon_0)\mu}.$$

We define $r^{(t)} = \nabla F(x^{(t)}) - H^{(t)} p^{(t)}$. Then we have that the direction vector satisfies

$$\|p^{(t)}\| = \|[H^{(t)}]^{-1}\| (\|r^{(t)}\| + \|\nabla F(x^{(t)})\|) \leq \frac{2}{(1 - \epsilon_0)\mu} \|\nabla F(x^{(t)})\|, \quad (16)$$

where the second inequality is because

$$\|r^{(t)}\| = \|\nabla F(x^{(t)}) - H^{(t)} p^{(t)}\| \leq \frac{\epsilon_1}{\kappa} \|\nabla F(x^{(t)})\| \leq \|\nabla F(x^{(t)})\|.$$

Hence, with $\|x - x^*\| \leq \gamma$, combining condition (15), we have

$$\begin{aligned}
& \|[\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}\| \int_0^1 \|\nabla^2 F(x^{(t)} + sp^{(t)}) - \nabla^2 F(x^{(t)})\| \|p^{(t)}\| ds \\
& \leq \|[\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}\| \int_0^1 \frac{\mu\eta(t)}{\sqrt{\kappa}} \|p^{(t)}\| ds \\
& \leq \|[\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}\| \frac{2}{(1 - \epsilon_0)\mu} \frac{\mu\eta(t)}{\sqrt{\kappa}} \|\nabla F(x^{(t)})\| \\
& \leq \frac{2\eta(t)}{1 - \epsilon_0} \frac{\|[\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}\|}{\sqrt{\kappa}\lambda_{\min}([\nabla^2 F(x^{(t)})]^{-\frac{1}{2}})} \lambda_{\min}([\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}) \|\nabla F(x^{(t)})\| \\
& \leq \frac{2\eta(t)}{1 - \epsilon_0} \|\nabla F(x^{(t)})\|_M.
\end{aligned}$$

Therefore, we have

$$\begin{aligned}
\|\nabla F(x^{(t+1)})\|_M & \leq \epsilon_0 \|\nabla F(x^{(t)})\|_M + \frac{\epsilon_1}{1 - \epsilon_0} \|\nabla F(x^{(t)})\|_M + \frac{2\eta(t)}{1 - \epsilon_0} \|\nabla F(x^{(t)})\|_M \\
& = \left(\epsilon_0 + \frac{\epsilon_1}{1 - \epsilon_0} + \frac{2\eta(t)}{1 - \epsilon_0} \right) \|\nabla F(x^{(t)})\|_M.
\end{aligned}$$

Now, we show the relationship between $\|\cdot\|_M$ and $\|\cdot\|_{M^*}$. By Eqn. (14), we have

$$-\frac{\nu(t)}{\lambda_{\max}(\nabla^2 F(x^*))} u^T u \leq u^T ([\nabla^2 F(x^*)]^{-1} - [\nabla^2 F(x^{(t)})]^{-1}) u \leq \frac{\nu(t)}{\lambda_{\max}(\nabla^2 F(x^*))} u^T u,$$

for any nonzero $u \in \mathbb{R}^d$, which implies that

$$(1 - \nu(t)) u^T [\nabla^2 F(x^{(t)})]^{-1} u \leq u^T [\nabla^2 F(x^*)]^{-1} u \leq (1 + \nu(t)) u^T [\nabla^2 F(x^{(t)})]^{-1} u.$$

That is,

$$(1 - \nu(t)) \|u\|_M \leq \|u\|_{M^*} \leq (1 + \nu(t)) \|u\|_M.$$

By this relationship between $\|\cdot\|_M$ and $\|\cdot\|_{M^*}$, we get

$$\|\nabla F(x^{(t+1)})\|_{M^*} \leq \left(\epsilon_0 + \frac{\epsilon_1}{1 - \epsilon_0} + \frac{2\eta(t)}{1 - \epsilon_0} \right) \frac{1 + \nu(t)}{1 - \nu(t)} \|\nabla F(x^{(t)})\|_{M^*}.$$

Second, we consider the case that $\nabla^2 F(x)$ is Lipschitz continuous with parameter \hat{L} . We have that the direction vector satisfies

$$\|p^{(t)}\| \leq \frac{2}{(1 - \epsilon_0)\lambda_{\min}(\nabla^2 F(x^{(t)}))} \|\nabla F(x^{(t)})\|.$$

Because $\nabla^2 F(x)$ is Lipschitz continuous with parameter \hat{L} , we have

$$\begin{aligned}
& \|[\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}\| \int_0^1 \|\nabla^2 F(x^{(t)} + sp^{(t)}) - \nabla^2 F(x^{(t)})\| \|p^{(t)}\| ds \\
& \leq \|[\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}\| \int_0^1 s \hat{L} \|p^{(t)}\|^2 ds \\
& = \frac{\hat{L}}{2} \|[\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}\| \lambda_{\min}^{-2}([\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}) \lambda_{\min}^2([\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}) \|p^{(t)}\|^2 \\
& \leq \frac{\hat{L}}{2} \|[\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}\| \lambda_{\min}^{-2}([\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}) \left(\frac{2}{(1 - \epsilon_0) \lambda_{\min}(\nabla^2 F(x^{(t)}))} \right)^2 \|\nabla F(x^{(t)})\|_M^2 \\
& = \frac{2\hat{L}\lambda_{\max}(\nabla^2 F(x^{(t)}))}{(1 - \epsilon_0)^2 \lambda_{\min}^2(\nabla^2 F(x^{(t)})) \sqrt{\lambda_{\min}(\nabla^2 F(x^{(t)}))}} \|\nabla F(x^{(t)})\|_M^2 \\
& \leq \frac{2}{(1 - \epsilon_0)^2} \cdot \frac{\hat{L}\kappa}{\mu\sqrt{\mu}} \|\nabla F(x^{(t)})\|_M^2.
\end{aligned}$$

Thus, we have

$$\|\nabla F(x^{(t+1)})\|_M \leq \left(\epsilon_0 + \frac{\epsilon_1}{1 - \epsilon_0} \right) \|\nabla F(x^{(t)})\|_M + \frac{2}{(1 - \epsilon_0)^2} \cdot \frac{\hat{L}\kappa}{\mu\sqrt{\mu}} \|\nabla F(x^{(t)})\|_M^2.$$

By the Lipschitz continuity of $\nabla^2 F(x)$ and the condition

$$\|x^{(t)} - x^*\| \leq \frac{\mu}{\hat{L}\kappa} \leq \frac{\lambda_{\min}(\nabla^2 F(x^*))}{\hat{L}\kappa(\nabla^2 F(x^{(t)}))},$$

we obtain

$$\begin{aligned}
\|[\nabla^2 F(x^*)]^{-1} - [\nabla^2 F(x^{(t)})]^{-1}\| & \leq \|[\nabla^2 F(x^*)]^{-1}\| \|[\nabla^2 F(x^{(t)})]^{-1}\| \|\nabla^2 F(x^*) - \nabla^2 F(x^{(t)})\| \\
& \leq \hat{L} \|[\nabla^2 F(x^*)]^{-1}\| \|[\nabla^2 F(x^{(t)})]^{-1}\| \|x^{(t)} - x^*\| \\
& \leq \nu(t) \lambda_{\min}([\nabla^2 F(x^{(t)})]^{-1}).
\end{aligned}$$

Hence, we can obtain that for any $u \in \mathbb{R}^d$,

$$-\nu(t) \lambda_{\min}([\nabla^2 F(x^{(t)})]^{-1}) u^T y \leq y^T ([\nabla^2 F(x^*)]^{-1} - [\nabla^2 F(x^{(t)})]^{-1}) y \leq \nu(t) \lambda_{\min}([\nabla^2 F(x^{(t)})]^{-1}) y^T y,$$

which yields

$$(1 - \nu(t)) u^T [\nabla^2 F(x^{(t)})]^{-1} u \leq u^T [\nabla^2 F(x^*)]^{-1} u \leq (1 + \nu(t)) u^T [\nabla^2 F(x^{(t)})]^{-1} u.$$

That is,

$$(1 - \nu(t)) \|u\|_M \leq \|u\|_{M^*} \leq (1 + \nu(t)) \|u\|_M.$$

Accordingly, we have

$$\|\nabla F(x^{(t+1)})\|_{M^*} \leq \left(\epsilon_0 + \frac{\epsilon_1}{1 - \epsilon_0} \right) \frac{1 + \nu(t)}{1 - \nu(t)} \|\nabla F(x^{(t)})\|_{M^*} + \frac{2}{(1 - \epsilon_0)^2} \cdot \frac{\hat{L}\kappa}{\mu\sqrt{\mu}} \frac{(1 + \nu(t))^2}{1 - \nu(t)} \|\nabla F(x^{(t)})\|_{M^*}^2.$$

■

Appendix C. Proofs of theorems of Section 4

Proof of Theorem 4 If S is an ϵ_0 -subspace embedding matrix w.r.t. $B(x^{(t)})$, then we have

$$(1 - \epsilon_0)\nabla^2 F(x^{(t)}) \preceq [B(x^{(t)})]^T S^T S B(x^{(t)}) \preceq (1 + \epsilon_0)\nabla^2 F(x^{(t)}). \quad (17)$$

By simple transformation and omitting ϵ_0^2 , (17) can be transformed into

$$(1 - \epsilon_0)[B(x^{(t)})]^T S^T S \nabla^2 B(x^{(t)}) \preceq \nabla^2 F(x^{(t)}) \preceq (1 + \epsilon_0)[B(x^{(t)})]^T S^T S B(x^{(t)}).$$

The convergence rate can be derived directly from Theorem 3. ■

Proof of Corollary 5 If $\nabla^2 F(x)$ is not Lipschitz continuous, then we have

$$\begin{aligned} \limsup_{t \rightarrow \infty} \frac{\|\nabla F(x^{(t+1)})\|_{M^*}}{\|\nabla F(x^{(t)})\|_{M^*}} &= \limsup_{t \rightarrow \infty} [\epsilon_0(t) + \eta(t)/(1 - \epsilon_0(t))] \frac{1 + \nu(t)}{1 - \nu(t)} \\ &= \limsup_{t \rightarrow \infty} \left(\frac{1}{\log(1+t)} + \frac{\eta(t)}{1 - \log(1+t)} \right) \frac{1 + \nu(t)}{1 - \nu(t)} \\ &= 0, \end{aligned}$$

where $\eta(t) \rightarrow 0$ is because $\|\nabla^2 F(x^{(t)}) - \nabla^2 F(x^*)\| \rightarrow 0$ as $x^{(t)}$ approaches x^* .

If $\nabla^2 F(x)$ is Lipschitz continuous, then we have

$$\begin{aligned} \limsup_{t \rightarrow \infty} \frac{\|\nabla F(x^{(t+1)})\|_{M^*}}{\|\nabla F(x^{(t)})\|_{M^*}} &= \limsup_{t \rightarrow \infty} \left(\epsilon_0(t) \frac{1 + \nu(t)}{1 - \nu(t)} + \frac{2}{(1 - \epsilon_0(t))^2} \frac{\hat{L}\kappa}{\mu\sqrt{\mu}} \frac{(1 + \nu(t))^2}{1 - \nu(t)} \|\nabla F(x^{(t)})\|_{M^*} \right) \\ &= \limsup_{t \rightarrow \infty} \left(\frac{1}{\log(1+t)} \frac{1 + \nu(t)}{1 - \nu(t)} + \frac{2}{(1 - \log(1+t))^2} \frac{\hat{L}\kappa}{\mu\sqrt{\mu}} \frac{(1 + \nu(t))^2}{1 - \nu(t)} \|\nabla F(x^{(t)})\|_{M^*} \right) \\ &= 0. \end{aligned}$$
■

Appendix D. Proofs of theorems of Section 5

Proof of Theorem 6 By Lemma 14, when $|\mathcal{S}| \geq \frac{16K^2 \log(2d/\delta)}{\sigma^2 \epsilon_0^2}$, $H^{(t)}$ has the following property:

$$\|H^{(t)} - \nabla^2 F(x^{(t)})\| \leq \epsilon_0 \sigma.$$

The above property implies the following:

$$\begin{aligned} |y^T (H^{(t)} - \nabla^2 F(x^{(t)})) y| &\leq \epsilon_0 \sigma y^T y, \\ \Rightarrow -\epsilon_0 \sigma y^T y &\leq y^T (H^{(t)} - \nabla^2 F(x^{(t)})) y \leq \epsilon_0 \sigma y^T y \\ \Rightarrow H^{(t)} - \epsilon_0 \sigma I &\preceq \nabla^2 F(x^{(t)}) \preceq H^{(t)} + \epsilon_0 \sigma I \\ \Rightarrow (1 - \epsilon_0) H^{(t)} &\preceq \nabla^2 F(x^{(t)}) \preceq (1 + \epsilon_0) H^{(t)}. \end{aligned}$$

The convergence rate can be derived directly from Theorem 3. ■

Proof of Theorem 7

By Lemma 14, when $|\mathcal{S}| \geq \frac{16K^2 \log(2d/\delta)}{\beta^2}$, we have

$$\|\nabla^2 F(x^{(t)}) - H_{|\mathcal{S}|}^{(t)}\| \leq \beta,$$

with probability at least $1 - \delta$. Hence, we can derive

$$\begin{aligned} & |y^T (\nabla^2 F(x^{(t)}) - H_{|\mathcal{S}|}^{(t)}) y| \leq \beta y^T y \\ \Rightarrow & y^T H_{|\mathcal{S}|}^{(t)} y - \beta y^T y \leq y^T \nabla^2 F(x^{(t)}) y \leq y^T H_{|\mathcal{S}|}^{(t)} y + \beta y^T y \\ \Rightarrow & y^T H^{(t)} y - \alpha y^T y - \beta y^T y \leq y^T \nabla^2 F(x^{(t)}) y \leq y^T H^{(t)} y - \alpha y^T y + \beta y^T y \\ \Rightarrow & y^T H^{(t)} y - (\alpha + \beta) y^T y \stackrel{(1)}{\leq} y^T \nabla^2 F(x^{(t)}) y \stackrel{(2)}{\leq} y^T H^{(t)} y + (\beta - \alpha) y^T y. \end{aligned}$$

Now we first consider $\stackrel{(1)}{\leq}$ case, we have

$$\begin{aligned} & y^T H^{(t)} y - (\alpha + \beta) y^T y \leq y^T \nabla^2 F(x^{(t)}) y \\ \Rightarrow & y^T H^{(t)} y \leq y^T \nabla^2 F(x^{(t)}) y + (\alpha + \beta) y^T y \\ \Rightarrow & y^T H^{(t)} y \leq y^T \nabla^2 F(x^{(t)}) y + \frac{\alpha + \beta}{\sigma} y^T \nabla^2 F(x^{(t)}) y \\ \Rightarrow & y^T H^{(t)} y \leq \left(1 + \frac{\alpha + \beta}{\sigma}\right) y^T \nabla^2 F(x^{(t)}) y \\ \Rightarrow & \left(1 - \frac{\alpha + \beta}{\sigma + \alpha + \beta}\right) y^T H^{(t)} y \leq y^T \nabla^2 F(x^{(t)}) y \\ \Rightarrow & \left(1 - \frac{\alpha + \beta}{\sigma + \alpha + \beta}\right) H^{(t)} \preceq F(x^{(t)}). \end{aligned}$$

For $\stackrel{(2)}{\leq}$ case, we consider two cases respectively. The first case is $\beta - \sigma/2 \leq \alpha \leq \beta$, and we have

$$\begin{aligned} & y^T \nabla^2 F(x^{(t)}) y \leq y^T H^{(t)} y + (\beta - \alpha) y^T y \\ \Rightarrow & y^T \nabla^2 F(x^{(t)}) y - (\beta - \alpha) y^T y \leq y^T H^{(t)} y \\ \Rightarrow & y^T \nabla^2 F(x^{(t)}) y - \frac{\beta - \alpha}{\sigma} y^T \nabla^2 F(x^{(t)}) y \leq y^T H^{(t)} y \\ \Rightarrow & \left(1 - \frac{\beta - \alpha}{\sigma}\right) y^T \nabla^2 F(x^{(t)}) y \leq y^T H^{(t)} y \\ \Rightarrow & y^T \nabla^2 F(x^{(t)}) y \leq \left(1 + \frac{\beta - \alpha}{\sigma - (\beta - \alpha)}\right) y^T H^{(t)} y \\ \Rightarrow & \nabla^2 F(x^{(t)}) \preceq \left(1 + \frac{\beta - \alpha}{\sigma + \alpha - \beta}\right) H^{(t)}. \end{aligned}$$

For the case $\beta < \alpha$, we can derive

$$\begin{aligned} y^T \nabla^2 F(x^{(t)}) y &\leq y^T H^{(t)} y + (\beta - \alpha) y^T y \leq y^T H^{(t)} y \\ \Rightarrow \nabla^2 F(x^{(t)}) &\preceq (1 + 0) H^{(t)}. \end{aligned}$$

Hence, for $\beta - \sigma \leq \alpha$, we have

$$\left(1 - \frac{\alpha + \beta}{\sigma + \alpha + \beta}\right) H^{(t)} \preceq F(x^{(t)}) \preceq \left(1 + \frac{\beta - \alpha}{\sigma + \alpha - \beta}\right) H^{(t)}.$$

Therefore, ϵ_0 in Theorem 3 can be set as follows:

$$\epsilon_0 = \max \left(\frac{\beta - \alpha}{\sigma + \alpha - \beta}, \frac{\alpha + \beta}{\sigma + \alpha + \beta} \right).$$

The convergence properties can be derived from Theorem 3 directly. ■

Proof of Theorem 8

We denote the SVD of $H_S^{(t)}$ as follows

$$H_S^{(t)} = U \hat{\Lambda} U^T = U_r \hat{\Lambda}_r U_r^T + U_{\setminus r} \hat{\Lambda}_{\setminus r} U_{\setminus r}^T.$$

By Lemma 14, when $|\mathcal{S}| \geq \frac{16K^2 \log(2d/\delta)}{\beta^2}$, we have

$$\|\nabla^2 F(x^{(t)}) - H_{|\mathcal{S}|}^{(t)}\| \leq \beta,$$

with probability at least $1 - \delta$. Hence, we can derive

$$\begin{aligned} &|y^T (\nabla^2 F(x^{(t)}) - H_{|\mathcal{S}|}^{(t)}) y| \leq \beta y^T y \\ \Rightarrow y^T H_{|\mathcal{S}|}^{(t)} y - \beta y^T y &\leq y^T \nabla^2 F(x^{(t)}) y \leq y^T H_{|\mathcal{S}|}^{(t)} y + \beta y^T y \\ \Rightarrow y^T H^{(t)} y + y^T U_{\setminus r} (\hat{\Lambda}_{\setminus r} - \hat{\lambda}_{r+1}^{(t)} I) U_{\setminus r}^T y - \beta y^T y &\leq y^T \nabla^2 F(x^{(t)}) y \\ &\leq y^T H^{(t)} y + y^T U_{\setminus r} (\hat{\Lambda}_{\setminus r} - \hat{\lambda}_{r+1}^{(t)} I) U_{\setminus r}^T y + \beta y^T y \\ \Rightarrow y^T H^{(t)} y - y^T U \begin{bmatrix} \beta I_r & \\ & (\beta + \hat{\lambda}_{r+1}^{(t)}) I_{\setminus r} - \hat{\Lambda}_{\setminus r} \end{bmatrix} U^T y &\stackrel{(1)}{\leq} y^T \nabla^2 F(x^{(t)}) y \\ &\stackrel{(2)}{\leq} y^T H^{(t)} y + y^T U \begin{bmatrix} \beta I_r & \\ & (\beta - \hat{\lambda}_{r+1}^{(t)}) I_{\setminus r} + \hat{\Lambda}_{\setminus r} \end{bmatrix} U^T y \end{aligned}$$

Now we first consider $\stackrel{(1)}{\leq}$ case, we have

$$\begin{aligned}
& y^T H^{(t)} y - y^T U \begin{bmatrix} \beta I_r & \\ & (\beta + \hat{\lambda}_{r+1}^{(t)}) I_{\setminus r} - \hat{\Lambda}_{\setminus r} \end{bmatrix} U^T y \stackrel{(1)}{\leq} y^T \nabla^2 F(x^{(t)}) y \\
& \Rightarrow y^T H^{(t)} y \leq y^T \nabla^2 F(x^{(t)}) y + (\beta + \hat{\lambda}_{r+1}^{(t)}) y^t y \\
& \Rightarrow y^T H^{(t)} y \leq y^T \nabla^2 F(x^{(t)}) y + \frac{\beta + \hat{\lambda}_{r+1}^{(t)}}{\sigma} y^T \nabla^2 F(x^{(t)}) y \\
& \Rightarrow y^T H^{(t)} y \leq y^T \nabla^2 F(x^{(t)}) y + \frac{2\beta + \hat{\lambda}_{r+1}}{\sigma} y^T \nabla^2 F(x^{(t)}) y \\
& \Rightarrow \left(1 - \frac{2\beta + \lambda_{r+1}^{(t)}}{\sigma + 2\beta + \lambda_{r+1}^{(t)}} \right) y^T H^{(t)} y \leq y^T \nabla^2 F(x^{(t)}) y.
\end{aligned}$$

Hence we have

$$\left(1 + \frac{\beta}{\lambda_{r+1}^{(t)} - \beta} \right) H^{(t)} \preceq \nabla^2 F(x).$$

Now we first consider $\stackrel{(2)}{\leq}$ case, we have

$$\begin{aligned}
y^T \nabla^2 F(x^{(t)}) y & \leq y^T H^{(t)} y + y^T U \begin{bmatrix} \beta I_r & \\ & (\beta - \hat{\lambda}_{r+1}^{(t)}) I_{\setminus r} + \hat{\Lambda}_{\setminus r} \end{bmatrix} U^T y \\
& \leq y^T H^{(t)} y + \frac{\beta}{\hat{\lambda}_{r+1}^{(t)}} y^T H^{(t)} y \\
& \leq \left(1 + \frac{\beta}{\lambda_{r+1}^{(t)} - \beta} \right) y^T H^{(t)} y,
\end{aligned}$$

where the last inequality is because $\lambda_{r+1}^{(t)} - \beta \leq \hat{\lambda}_{r+1}^{(t)}$. Hence, we have

$$\nabla^2 F(x) \preceq \left(1 + \frac{\beta}{\lambda_{r+1}^{(t)} - \beta} \right) H^{(t)}.$$

Hence, we have

$$\epsilon_0 = \max \left(\frac{\beta}{\lambda_{r+1}^{(t)} - \beta}, \frac{2\beta + \lambda_{r+1}^{(t)}}{\sigma + 2\beta + \lambda_{r+1}^{(t)}} \right) < 1,$$

because $\beta \leq \frac{\lambda_{r+1}^{(t)}}{2}$.

The convergence properties can be derived directly by Theorem 3. ■

The proof of Theorem 9 is almost the same with Theorem 3. For completeness, we give the detailed proof as follows.

Proof of Theorem 9

By Taylor's theorem, we obtain

$$\begin{aligned}
& \nabla F(x^{(t+1)}) \\
&= \nabla F(x^{(t)}) + \nabla^2 F(x^{(t)})(-p^{(t)}) + \int_0^1 [\nabla^2 F(x^{(t)} + sp^{(t)}) - \nabla^2 F(x^{(t)})](-p^{(t)})ds \\
&= \nabla F(x^{(t)}) - \nabla^2 F(x^{(t)})[H^{(t)}]^{-1}\nabla F(x^{(t)}) + \nabla^2 F(x^{(t)})[H^{(t)}]^{-1}\nabla F(x^{(t)}) - \nabla^2 F(x^{(t)})p^{(t)} \\
&\quad + \int_0^1 [\nabla^2 F(x^{(t)} + sp^{(t)}) - \nabla^2 F(x^{(t)})](-p^{(t)})ds \\
&= [\nabla^2 F(x^{(t)})]^{\frac{1}{2}} \left(I - [\nabla^2 F(x^{(t)})]^{\frac{1}{2}}[H^{(t)}]^{-1}[\nabla^2 F(x^{(t)})]^{\frac{1}{2}} \right) [\nabla^2 F(x^{(t)})]^{-\frac{1}{2}} \nabla F(x^{(t)}) \\
&\quad + \nabla^2 F(x^{(t)})([H^{(t)}]^{-1}\nabla F(x^{(t)}) - p^{(t)}) + \int_0^1 [\nabla^2 F(x^{(t)} + sp^{(t)}) - \nabla^2 F(x^{(t)})](-p^{(t)})ds.
\end{aligned}$$

Hence, we have the following identity

$$\begin{aligned}
[\nabla^2 F(x^{(t)})]^{-\frac{1}{2}} \nabla F(x^{(t+1)}) &= \left(I - [\nabla^2 F(x^{(t)})]^{\frac{1}{2}}[H^{(t)}]^{-1}[\nabla^2 F(x^{(t)})]^{\frac{1}{2}} \right) [\nabla^2 F(x^{(t)})]^{-\frac{1}{2}} \nabla F(x^{(t)}) \\
&\quad + [\nabla^2 F(x^{(t)})]^{\frac{1}{2}}([H^{(t)}]^{-1}\nabla F(x^{(t)}) - p^{(t)}) \\
&\quad + [\nabla^2 F(x^{(t)})]^{-\frac{1}{2}} \int_0^1 [\nabla^2 F(x^{(t)} + sp^{(t)}) - \nabla^2 F(x^{(t)})](-p^{(t)})ds.
\end{aligned}$$

Further more, we define $M = [\nabla^2 F(x^{(t)})]^{-1}$, we can obtain

$$\begin{aligned}
\|\nabla F(x^{(t+1)})\|_M &\leq \left\| I - [\nabla^2 F(x^{(t)})]^{\frac{1}{2}}[H^{(t)}]^{-1}[\nabla^2 F(x^{(t)})]^{\frac{1}{2}} \right\| \|\nabla F(x^{(t)})\|_M \\
&\quad + \|[\nabla^2 F(x^{(t)})]^{\frac{1}{2}}\| \| [H^{(t)}]^{-1}(\nabla F(x^{(t)}) - g(x^{(t)})) \| \\
&\quad + \|[\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}\| \int_0^1 \|\nabla^2 F(x^{(t)} + sp^{(t)}) - \nabla^2 F(x^{(t)})\| \|p^{(t)}\| ds.
\end{aligned}$$

We will bound the three terms on the right hand of above equation seperately.

For the first term, using Lemma 11, we have

$$\left\| I - [\nabla^2 F(x^{(t)})]^{\frac{1}{2}}[H^{(t)}]^{-1}[\nabla^2 F(x^{(t)})]^{\frac{1}{2}} \right\| \|\nabla F(x^{(t)})\|_M \leq \epsilon_0 \|\nabla F(x^{(t)})\|_M.$$

For the second term, by the fact that $\|AB\| \geq \|A\|\sigma_{\min}(B)$ and condition $\|g(x^{(t)}) - \nabla F(x^{(t)})\| \leq \epsilon_2 \|\nabla F(x^{(t)})\|$, we obtain

$$\begin{aligned}
& \|[\nabla^2 F(x^{(t)})]^{\frac{1}{2}}\| \| [H^{(t)}]^{-1}(\nabla F(x^{(t)}) - g(x^{(t)})) \| \\
&= \frac{\|[\nabla^2 F(x^{(t)})]^{\frac{1}{2}}\|}{\lambda_{\min}([\nabla^2 F(x^{(t)})]^{-\frac{1}{2}})} \lambda_{\min}([\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}) \| [H^{(t)}]^{-1} \| \|\nabla F(x^{(t)}) - g(x^{(t)})\| \\
&\leq \epsilon_2 \frac{\|[\nabla^2 F(x^{(t)})]^{\frac{1}{2}}\|}{\lambda_{\min}([\nabla^2 F(x^{(t)})]^{-\frac{1}{2}})} \| [H^{(t)}]^{-1} \| \|\nabla F(x^{(t)})\|_M \\
&\leq \frac{\epsilon_2}{1 - \epsilon_0} \|\nabla F(x^{(t)})\|_M.
\end{aligned}$$

For the third term, we bound it for the case that $\nabla^2 F(x)$ is not Lipschitz continuous and the case $\nabla^2 F(x)$ is Lipschitz continuous respectively.

(a) Now we consider the case that $\nabla^2 F(x)$ is not Lipschitz continuous but is continuous close to the optimal point x^* . Because $\nabla^2 F(x)$ is continuous near x^* , there exists a sufficient small value γ such that Eqn. (14) and Eqn. (15) hold when $\|x^{(t)} - x^*\| \leq \gamma$.

By μ -strong convexity, we have $\|[\nabla^2 F(x^t)]^{-1}\| \leq \frac{1}{\mu}$ for all $x^{(t)}$ sufficiently close to x^* . Then we have

$$\|p^{(t)}\| = \|[H^{(t)}]^{-1}\| \|g(x^{(t)})\| \leq \frac{1 + \epsilon_2/\kappa}{(1 - \epsilon_0)\mu} \|\nabla F(x^{(t)})\| \leq \frac{2}{(1 - \epsilon_0)\mu} \|\nabla F(x^{(t)})\|.$$

Hence, with $\|x - x^*\| \leq \delta$, combining condition (15), we have

$$\begin{aligned} & \|[\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}\| \int_0^1 \|\nabla^2 F(x^{(t)} + sp^{(t)}) - \nabla^2 F(x^{(t)})\| \|p^{(t)}\| ds \\ & \leq \|[\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}\| \int_0^1 \frac{\mu\eta(t)}{\sqrt{\kappa}} \|p^{(t)}\| ds \\ & \leq \|[\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}\| \frac{2}{(1 - \epsilon_0)\mu} \frac{\mu\eta(t)}{\sqrt{\kappa}} \|\nabla F(x^{(t)})\| \\ & \leq \frac{2\eta(t)}{1 - \epsilon_0} \frac{\|[\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}\|}{\sqrt{\kappa}\lambda_{\min}([\nabla^2 F(x^{(t)})]^{-\frac{1}{2}})} \lambda_{\min}([\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}) \|\nabla F(x^{(t)})\| \\ & \leq \frac{2\eta(t)}{1 - \epsilon_0} \|\nabla F(x^{(t)})\|_M, \end{aligned}$$

Therefore, we have

$$\begin{aligned} \|\nabla F(x^{(t+1)})\|_M & \leq \epsilon_0 \|\nabla F(x^{(t)})\|_M + \frac{\epsilon_2}{1 - \epsilon_0} \|\nabla F(x^{(t)})\|_M + \frac{2\eta(t)}{1 - \epsilon_0} \|\nabla F(x^{(t)})\|_M \\ & = \left(\epsilon_0 + \frac{\epsilon_2}{1 - \epsilon_0} + \frac{2\eta(t)}{1 - \epsilon_0} \right) \|\nabla F(x^{(t)})\|_M. \end{aligned}$$

By Eqn. (14), we have

$$\begin{aligned} & -\frac{\nu(t)}{\lambda_{\max}(\nabla^2 F(x^*))} y^T y \leq y^T ([\nabla^2 F(x^*)]^{-1} - [\nabla^2 F(x^{(t)})]^{-1}) y \leq \frac{\nu(t)}{\lambda_{\max}(\nabla^2 F(x^*))} y^T y, \\ \Rightarrow & (1 - \nu(t)) y^T [\nabla^2 F(x^{(t)})]^{-1} y \leq y^T [\nabla^2 F(x^*)]^{-1} y \leq (1 + \nu(t)) y^T [\nabla^2 F(x^{(t)})]^{-1} y \\ \Rightarrow & (1 - \nu(t)) \|y\|_M \leq \|y\|_{M^*} \leq (1 + \nu(t)) \|y\|_M. \end{aligned}$$

By this relationship between $\|\cdot\|_M$ and $\|\cdot\|_{M^*}$, we get

$$\|\nabla F(x^{(t+1)})\|_{M^*} \leq \left(\epsilon_0 + \frac{\epsilon_2}{1 - \epsilon_0} + \frac{2\eta(t)}{1 - \epsilon_0} \right) \frac{1 + \nu(t)}{1 - \nu(t)} \|\nabla F(x^{(t)})\|_{M^*}.$$

(b) Now we consider the case that $\nabla^2 F(x)$ is Lipschitz continuous with parameter \hat{L} . The same to the previous proof, we have

$$\|p^{(t)}\| = \|[H^{(t)}]^{-1}\| \|g(x^{(t)})\| \leq \frac{1 + \epsilon_2/\kappa}{(1 - \epsilon_0)\mu} \|\nabla F(x^{(t)})\| \leq \frac{2}{(1 - \epsilon_0)\mu} \|\nabla F(x^{(t)})\|.$$

Because $\nabla^2 F(x)$ is Lipschitz continuous with parameter \hat{L} , we have

$$\begin{aligned}
& \|[\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}\| \int_0^1 \|\nabla^2 F(x^{(t)} + sp^{(t)}) - \nabla^2 F(x^{(t)})\| \|p^{(t)}\| ds \\
& \leq \|[\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}\| \int_0^1 sL \|p^{(t)}\|^2 ds \\
& = \frac{\hat{L}}{2} \|[\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}\| \lambda_{\min}^{-2}([\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}) \lambda_{\min}^2([\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}) \|p^{(t)}\|^2 \\
& \leq \frac{\hat{L}}{2} \|[\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}\| \lambda_{\min}^{-2}([\nabla^2 F(x^{(t)})]^{-\frac{1}{2}}) \left(\frac{2}{(1 - \epsilon_0) \lambda_{\min}(\nabla^2 F(x^{(t)}))} \right)^2 \|\nabla F(x^{(t)})\|_M^2 \\
& = \frac{2\hat{L}\lambda_{\max}(\nabla^2 F(x^{(t)}))}{(1 - \epsilon_0)^2 \lambda_{\min}^2(\nabla^2 F(x^{(t)})) \sqrt{\lambda_{\min}(\nabla^2 F(x^{(t)}))}} \|\nabla F(x^{(t)})\|_M^2 \\
& \leq \frac{2\hat{L}\kappa}{(1 - \epsilon_0)^2 \mu \sqrt{\mu}} \|\nabla F(x^{(t)})\|_M^2.
\end{aligned}$$

Hence, we have

$$\|\nabla F(x^{(t+1)})\|_M \leq \left(\epsilon_0 + \frac{\epsilon_2}{1 - \epsilon_0} \right) \|\nabla F(x^{(t)})\|_M + \frac{2\hat{L}\kappa}{(1 - \epsilon_0)^2 \mu \sqrt{\mu}} \|\nabla F(x^{(t)})\|_M^2.$$

By the Lipschitz continuity of $\nabla^2 F(x)$ and the condition

$$\|x^{(t)} - x^*\| \leq \frac{\mu}{\hat{L}\kappa} \leq \frac{\lambda_{\min}(\nabla^2 F(x^*))}{\hat{L}\kappa(\nabla^2 F(x^{(t)}))},$$

we obtain

$$\begin{aligned}
\|[\nabla^2 F(x^*)]^{-1} - [\nabla^2 F(x^{(t)})]^{-1}\| & \leq \|[\nabla^2 F(x^*)]^{-1}\| \|[\nabla^2 F(x^{(t)})]^{-1}\| \|\nabla^2 F(x^*) - \nabla^2 F(x^{(t)})\| \\
& \leq \hat{L} \|[\nabla^2 F(x^*)]^{-1}\| \|[\nabla^2 F(x^{(t)})]^{-1}\| \|x^{(t)} - x^*\| \\
& \leq \nu(t) \lambda_{\min}([\nabla^2 F(x^{(t)})]^{-1}).
\end{aligned}$$

Hence, we can derive

$$\begin{aligned}
& -\nu(t) \lambda_{\min}([\nabla^2 F(x^{(t)})]^{-1}) y^T y \leq y^T ([\nabla^2 F(x^*)]^{-1} - [\nabla^2 F(x^{(t)})]^{-1}) y \leq \nu(t) \lambda_{\min}([\nabla^2 F(x^{(t)})]^{-1}) y^T y, \\
& \Rightarrow (1 - \nu(t)) y^T [\nabla^2 F(x^{(t)})]^{-1} y \leq y^T [\nabla^2 F(x^*)]^{-1} y \leq (1 + \nu(t)) y^T [\nabla^2 F(x^{(t)})]^{-1} y \\
& \Rightarrow (1 - \nu(t)) \|y\|_M \leq \|y\|_{M^*} \leq (1 + \nu(t)) \|y\|_M.
\end{aligned}$$

Hence, we have

$$\|\nabla F(x^{(t+1)})\|_{M^*} \leq \left(\epsilon_0 + \frac{\epsilon_2}{1 - \epsilon_0} \right) \frac{1 + \nu(t)}{1 - \nu(t)} \|\nabla F(x^{(t)})\|_{M^*} + \frac{2\hat{L}\kappa}{(1 - \epsilon_0)^2 \mu \sqrt{\mu}} \frac{(1 + \nu(t))^2}{1 - \nu(t)} \|\nabla F(x^{(t)})\|_{M^*}^2.$$

■

References

- [1] Naman Agarwal, Brian Bullins, and Elad Hazan. Second order stochastic optimization in linear time. *arXiv preprint arXiv:1602.03943*, 2016.
- [2] Raghu Bollapragada, Richard Byrd, and Jorge Nocedal. Exact and inexact subsampled newton methods for optimization. *arXiv preprint arXiv:1609.08502*, 2016.
- [3] Richard H Byrd, Gillian M Chin, Will Neveitt, and Jorge Nocedal. On the use of stochastic hessian information in optimization methods for machine learning. *SIAM Journal on Optimization*, 21(3):977–995, 2011.
- [4] Kenneth L Clarkson and David P Woodruff. Low rank approximation and regression in input sparsity time. In *Proceedings of the forty-fifth annual ACM symposium on Theory of computing*, pages 81–90. ACM, 2013.
- [5] Andrew Cotter, Ohad Shamir, Nati Srebro, and Karthik Sridharan. Better mini-batch algorithms via accelerated gradient methods. In *Advances in neural information processing systems*, pages 1647–1655, 2011.
- [6] Petros Drineas, Michael W Mahoney, and S Muthukrishnan. Sampling algorithms for l_2 regression and applications. In *Proceedings of the seventeenth annual ACM-SIAM symposium on Discrete algorithm*, pages 1127–1136. Society for Industrial and Applied Mathematics, 2006.
- [7] Petros Drineas, Malik Magdon-Ismail, Michael W Mahoney, and David P Woodruff. Fast approximation of matrix coherence and statistical leverage. *Journal of Machine Learning Research*, 13(Dec):3475–3506, 2012.
- [8] Murat A Erdogdu and Andrea Montanari. Convergence rates of sub-sampled newton methods. In *Advances in Neural Information Processing Systems*, pages 3034–3042, 2015.
- [9] N Halko, P G Martinsson, and J A Tropp. Finding Structure with Randomness : Probabilistic Algorithms for Matrix Decompositions. *SIAM Review*, 53(2):217–288, 2011.
- [10] Rie Johnson and Tong Zhang. Accelerating stochastic gradient descent using predictive variance reduction. In *Advances in Neural Information Processing Systems*, pages 315–323, 2013.
- [11] William B. Johnson and Joram Lindenstrauss. Extensions of Lipschitz mappings into a Hilbert space. *Contemporary mathematics*, 26(189-206), 1984.
- [12] Mu Li, Tong Zhang, Yuqiang Chen, and Alexander J Smola. Efficient mini-batch training for stochastic optimization. In *Proceedings of the 20th ACM SIGKDD international conference on Knowledge discovery and data mining*, pages 661–670. ACM, 2014.
- [13] Xiangrui Meng and Michael W Mahoney. Low-distortion subspace embeddings in input-sparsity time and applications to robust linear regression. In *Proceedings of the forty-fifth annual ACM symposium on Theory of computing*, pages 91–100. ACM, 2013.

- [14] Jelani Nelson and Huy L Nguyễn. Osnap: Faster numerical linear algebra algorithms via sparser subspace embeddings. In *Foundations of Computer Science (FOCS), 2013 IEEE 54th Annual Symposium on*, pages 117–126. IEEE, 2013.
- [15] Arkadi Nemirovski, Anatoli Juditsky, Guanghui Lan, and Alexander Shapiro. Robust stochastic approximation approach to stochastic programming. *SIAM Journal on Optimization*, 19(4):1574–1609, 2009.
- [16] Jorge Nocedal and Stephen Wright. *Numerical optimization*. Springer Science & Business Media, 2006.
- [17] Mert Pilanci and Martin J Wainwright. Newton sketch: A linear-time optimization algorithm with linear-quadratic convergence. *arXiv preprint arXiv:1505.02250*, 2015.
- [18] Herbert Robbins and Sutton Monro. A stochastic approximation method. *The annals of mathematical statistics*, pages 400–407, 1951.
- [19] Farbod Roosta-Khorasani and Michael W Mahoney. Sub-sampled newton methods ii: Local convergence rates. *arXiv preprint arXiv:1601.04738*, 2016.
- [20] Nicolas L Roux, Mark Schmidt, and Francis R Bach. A stochastic gradient method with an exponential convergence rate for finite training sets. In *Advances in Neural Information Processing Systems*, pages 2663–2671, 2012.
- [21] Mark Schmidt, Nicolas Le Roux, and Francis Bach. Minimizing finite sums with the stochastic average gradient. *arXiv preprint arXiv:1309.2388*, 2013.
- [22] Joel A Tropp et al. An introduction to matrix concentration inequalities. *Foundations and Trends® in Machine Learning*, 8(1-2):1–230, 2015.
- [23] David P Woodruff. Sketching as a tool for numerical linear algebra. *Foundations and Trends® in Theoretical Computer Science*, 10(1–2):1–157, 2014.
- [24] Peng Xu, Jiyan Yang, Farbod Roosta-Khorasani, Christopher Ré, and Michael W Mahoney. Sub-sampled newton methods with non-uniform sampling. In *Advances in Neural Information Processing Systems*, pages 3000–3008, 2016.
- [25] Lijun Zhang, Mehrdad Mahdavi, and Rong Jin. Linear convergence with condition number independent access of full gradients. In *Advance in Neural Information Processing Systems 26 (NIPS)*, pages 980–988, 2013.